For Research Use Only

CoraLite® Plus 488-conjugated TDG Recombinant antibody

Catalog Number: CL488-83780-3



Purification Method:

493 nm / 522 nm

Positive Controls:

Basic Information

Catalog Number: GenBank Accession Number: CL488-83780-3

BC037557 Protein A purification GeneID (NCBI): CloneNo.:

Size: 1000 µg/ml 6996 240715E12 **UNIPROT ID:** Source:

Recommended Dilutions: Rabbit Q13569 IF/ICC 1:50-1:500

Full Name: Excitation/Emission maxima Isotype: thymine-DNA glycosylase wavelengths:

Calculated MW: Immunogen Catalog Number:

AG4190 410 aa, 46 kDa

Applications

Tested Applications:

IF/ICC IF/ICC: HepG2 cells,

Species Specificity:

human

Background Information

TDG belongs to the TDG/mug DNA glycosylase family. TDG corrects G/T mispairs to G/C pairs. It is capable of hydrolyzing the carbon-nitrogen bond between the sugar-phosphate backbone of the DNA and a mispaired thymine. In addition to the G/T, it can remove thymine also from C/T and T/T mispairs in the order G/T >> C/T > T/T. It has no detectable activity on apyrimidinic sites and does not catalyze the removal of thymine from A/T pairs or from $single-stranded\ DNA.\ It\ can\ also\ remove\ uracil\ and\ 5-bromouracil\ from\ mispairs\ with\ guanine.\ RNF4\ interacts\ with$ and requires the base excision repair enzymes TDG and APE1 for active demethylation (PMID:20696907). TDG is modified by SUMO-1 and SUMO-2/3. The molecular weight of non-modified TDG is 46 kDa and modified TDG is 55-

Storage

Storage:


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using CoraLite® Plus 488 TDG antibody (CL488-83780-3, Clone: 240715E12) at dilution of 1:200, CL594-Phalloidin (red).