For Research Use Only

CoraLite®594-conjugated Caspase 3/p17/p19 Monoclonal antibody



Purification Method:

CloneNo.:

IF 1:50-1:500

588 nm / 604 nm

2G4B2

Protein A purification

Recommended Dilutions:

Excitation/Emission maxima

Catalog Number: CL594-66470

Featured Product

Basic Information

Catalog Number: CL594-66470

Size: 1000 µg/ml Source: Mouse

lgG1 Immunogen Catalog Number:

AG25029

Isotype:

Tested Applications: Species Specificity:

Human, mouse, rat

GenBank Accession Number:

BC016926 GeneID (NCBI):

UNIPROT ID: P42574 Full Name:

Calculated MW: 277 aa, 32 kDa

caspase 3, apoptosis-related cysteine wavelengths: peptidase

Observed MW: 32-35 kDa, 19 kDa, 17 kDa

Positive Controls:

IF: HepG2 cells,

Background Information

Caspases, a family of endoproteases, are critical players in cell regulatory networks controlling inflammation and cell death. Initiator caspases (caspase-2, -8, -9, -10, -11, and -12) cleave and activate downstream effector caspases (caspase-3, -6, and -7), which in turn execute apoptosis by cleaving targeted cellular proteins. Caspase 3 (also named CPP32, SCA-1, and Apopain) proteolytically cleaves poly(ADP-ribose) polymerase (PARP) at the beginning of apoptosis. Caspase 3 plays a key role in the activation of sterol regulatory element binding proteins (SREBPs) between the basic helix-loop-helix leucine zipper domain and the membrane attachment domain. Caspase 3 can also form heterocomplex with other proteins and performs the molecular mass of 50-70 kDa. This antibody can recognize p17, p19 and p32 of Caspase 3.

Storage

Applications


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using CoraLite®594 Caspase 3/p17/p19 antibody (CL594-66470, Clone: 2G4B2) at dilution of 1:200.