For Research Use Only

CoraLite®594-conjugated Beta Actin Recombinant antibody

Catalog Number: CL594-81115



Basic Information

Catalog Number: GenBank Accession Number: CL594-81115 BC002409

Size: Genel D (NCBI):

 $\begin{array}{lll} \text{1000} \; \mu \, \text{g/ml} & \text{60} \\ \\ \text{Source:} & \text{UNIPROT ID:} \\ \\ \text{Rabbit} & \text{P60709} \\ \\ \text{Isotype:} & \text{Full Name:} \\ \\ \text{IgG} & \text{actin, beta} \\ \end{array}$

Immunogen Catalog Number:Calculated MW:AG14521375 aa, 42 kDa

Observed MW: 42 kDa Purification Method:

Protein A purification

CloneNo.: 4H1

Recommended Dilutions: IF/ICC 1:50-1:500

Excitation/Emission maxima wavelengths:

588 nm / 604 nm

Applications

Tested Applications:

IF/ICC

Species Specificity:

human, mouse, rat, pig, canine

Positive Controls:

IF/ICC : MDCK cells,

Background Information

Beta Actin, also named as ACTB and F-Actin, belongs to the actin family. Actins are highly conserved globular proteins that are involved in various types of cell motility and are ubiquitously expressed in all eukaryotic cells. At least six isoforms of actins are known in mammals and other vertebrates: alpha (ACTC1, cardiac muscle 1), alpha 1 (ACTA1, skeletal muscle) and 2 (ACTA2, aortic smooth muscle), beta (ACTB), gamma 1 (ACTG1) and 2 (ACTG2, enteric smooth muscle). Beta and gamma 1 are two non-muscle actin proteins. Most actins consist of 376aa, while ACTG2 (rich in muscles) has 375aa and ACTG1(found in non-muscle cells) has only 374aa. Beta actin has been widely used as the internal control in RT-PCR and Western Blotting as a 42-kDa protein. However, the 37-40, 31, 15 kDa cleaved fragment of beta actin can be generated during apoptosis process. This antibody was generated against N-terminal region of human beta actin protein and can cross-react with other actins. (9173887, 11217076, 10229193)

Storage

Storage:


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (-20°C Methanol) fixed MDCK cells using CoraLite®594 Beta Actin antibody (CL594-81115, Clone: 4H1) at dilution of 1:200.