For Research Use Only

CoraLite®594 Anti-Human TLR3/CD283 Rabbit Recombinant Antibody



Catalog Number: CL594-98016

Basic Information

Catalog Number: GenBank Accession Number: CL594-98016 BC094737

Concentration: Genel D (NCBI): 100tests, 5 ul/test 7098

Source: UNIPROT ID: Rabbit 015455
Isotype: Full Name:

IgG toll-like receptor 3
Immunogen Catalog Number: Calculated MW:
AG34044 904 aa, 104 kDa

Purification Method: Protein A purification

CloneNo.: 230445F1

Excitation/Emission maxima

wavelengths: 588 nm / 604 nm

Applications

Tested Applications:

FC (Intra)

Species Specificity:

human

Background Information

TLR3 belongs to the Toll-like receptor family which is important in the innate immune response to pathogens. TLRs are highly conserved from Drosophila to humans and share structural and functional similarities. TLR3 is a nucleotide-sensing TLR that is activated by double-stranded RNA. Upon recognition of dsRNA, TLR3 transmits signals via the adaptor protein TICAM-1/TRIF, leading to the induction of type I IFN, cytokine/chemokine production, and dendritic cell (DC) maturation (PMID: 18262679).

Storage


Storage:

Store at 2-8 °C. Avoid exposure to light. Stable for one year after shipment.


Storage Buffer:

PBS with 0.09% sodium azide and 0.5% BSA

Selected Validation Data



1x10^6 human monocyte-derived immature dendritic cells were intracellularly stained with 5 ul Coralite®594 Anti-Human TLR3/CD283 Rabbit RecAb (CL594-98016, Clone: 230445F1) or Coralite®594 Rabbit IgG Isotype Control RecAb (CL594-98136, Clone: 240953C9). Cells were co-stained with Coralite® Plus 647 Anti-Human CD11c. Cells were incubated with FC Receptor Block prior to fixing and staining. Cells were fixed with 4% PFA and permeabilized with Flow



1x10^6 human monocyte-derived immature dendritic cells were intracellularly stained with 5 ul Coralite®594 Anti-Human TLR3/CD283 Rabbit RecAb (CL594-98016, Clone: 230445F1) (red), or Coralite®594 Rabbit IgG Isotype Control RecAb (CL594-98136, Clone: 240953C9) (blue). Cells were incubated with FC Receptor Block prior to fixing and staining. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).