For Research Use Only

CoraLite® Plus 647 Anti-Human ICAM-1/CD54 (15.2) Rabbit IgG Recombinant Antibody



Catalog Number: CL647-65509

Basic Information

Catalog Number: CL647-65509

Size:

100tests, 5 ul/test

Source: Rabbit

Isotype:

IgG

GenBank Accession Number:

BC015969 GeneID (NCBI):

3383

ENSEMBL Gene ID: ENSG00000090339

Full Name:

intercellular adhesion molecule 1

Calculated MW: 90 kDa

Purification Method:

Affinity purification CloneNo.:

Excitation/Emission maxima

wavelengths: 654 nm / 674 nm

Applications

Tested Applications:

FC

Species Specificity:

human

Background Information

ICAM-1 (CD54) is a 90-kDa transmembrane glycoprotein of the immunoglobulin superfamily and is critical for the firm attachment and transmigration of leukocytes out of blood vessels and into tissues (PMID: 19307690). ICAM-1 is expressed by several cell types, typically on endothelial cells and cells of the immune system, and its expression can be up-regulated by various stimuli, including TNF- α , IL-1 and thrombin (PMID: 3086451; 9694714; 15979056). It is a ligand for LFA-1 and Mac-1, serves as a receptor for rhinovirus, and is one of several receptors used by Plasmodium falciparum (PMID: 2566624; 2538244; 2475784).


Storage

Storage:


Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment. Storage Ruffer

PBS with 0.09% sodium azide and 0.5% BSA.

Selected Validation Data



1x10^6 human PBMCs were surface stained with 5 ul CoraLite® Plus 647 Anti-Human CD54 Rabbit Recombinant Antibody (CL647-65509, Clone:15.2) or CoraLite® Plus 647-conjugated Rabbit IgG control PolyAb (CL647-3000). Cells were not fixed. Cells were treated with FC Receptor Block prior to staining. Monocytes were gated.



1x10^6 human PBMCs were surface stained with 5 ul CoraLite® Plus 647 Anti-Human CD54 Rabbit Recombinant Antibody (CL647-65509, Clone:15.2) or CoraLite® Plus 647-conjugated Rabbit IgG control PolyAb (CL647-3000). Cells were not fixed. Cells were treated with FC Receptor Block prior to staining. Monocytes were gated.