For Research Use Only

CoraLite® Plus 647 Anti-Human CD86 (BU63) Mouse IgG2a Recombinant Antibody



Catalog Number: CL647-65592

Basic Information

Catalog Number:

CL647-65592

Size:

100tests, 5 ul/test

Source: Mouse

Isotype: lgG2a

GenBank Accession Number:

BC040261 GeneID (NCBI):

ENSEMBL Gene ID:

ENSG00000114013 Full Name:

CD86 molecule

Calculated MW: 329 aa, 38 kDa

Purification Method:

Protein A purification

CloneNo.: BU63

Excitation/Emission maxima

wavelengths: 654 nm / 674 nm

Applications

Tested Applications:

Species Specificity:

human

Background Information

CD86 (also known as B7.2) is a costimulatory molecule belonging to the immunoglobulin superfamily. Primarily expressed on antigen-presenting cells (APCs), including B cells, dendritic cells, and macrophages, CD86 is the ligand for two proteins at the cell surface of T cells, CD28 antigen and cytotoxic T-lymphocyte-associated protein 4. Binding of CD86 with CD28 antigen is a costimulatory signal for activation of the T-cell. Binding of CD86 with cytotoxic T $lymphocyte-associated\ protein\ 4\ negatively\ regulates\ T-cell\ activation\ and\ diminishes\ the\ immune\ response.$


Storage

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.


Storage Buffer:

PBS with 0.09% sodium azide.

Selected Validation Data



1x10^6 human PBMCs were surface stained with CoraLite® Plus 488 Anti-Human CD14 Rabbit Recombinant Antibody (CL488-98040, Clone: 230332D7) and 5 ul CoraLite® Plus 647 Anti-Human CD86 (BU63) Mouse IgG2a RecAb (CL647-65592, Clone: BU63) or CoraLite® Plus 647 Mouse IgG2a Isotype Control (C1.18.4) (CL647-65208, Clone: C1.18.4). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.



CL647-65592 CD86(BU63),APC-A

1x10^6 human PBMCs were surface stained with 5 ul CoraLite® Plus 647 Anti-Human CD86 (BU63) Mouse IgG2a RecAb (CL647-65592, Clone: BU63) (red) or CoraLite® Plus 647 Mouse IgG2a Isotype Control (C1.18.4) (CL647-65208, Clone: C1.18.4) (blue). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.