For Research Use Only

## CoraLite® Plus 647-conjugated DEK Monoclonal antibody



Catalog Number: CL647-66194

Featured Product

**Basic Information** 

Catalog Number: CL647-66194

Size: 1000 µg/ml

Source: Mouse Isotype: IgG2a

Immunogen Catalog Number:

AG8225

43 kDa Observed MW:

40-50 kDa

BC035259

7913

P35659

GeneID (NCBI):

**UNIPROT ID:** 

Full Name:

**DEK** oncogene Calculated MW:

GenBank Accession Number:

**Purification Method:** 

Protein A purification

CloneNo.: 5E3A6

Excitation/Emission maxima

wavelengths: 654 nm / 674 nm

**Applications** 

**Tested Applications:** 

FC (Intra)

Species Specificity: human, mouse, rat

**Background Information** 

Dek oncogene(DEK) is first recognized in a fusion with CAN nucleoporin protein, and it's a also autoantigen in patients with pquciarticular onset juvenile rheumatoid arthritis. As a site-specific binding protein, DEK may involve in transcriptional regulation and signal transdcution. Analysis by agarose gel celctrophoresis, DEK showed to reduce the efficiency with which chromation is replicated. It's one of the first components of the splicing complex, which remains associate with spliced exons dependent on prior splicing of pre-mRNA.


Storage

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



1x10^6 HeLa cells were intracellularly stained with 0.4 ug CoraLite® Plus 647-conjugated DEK Monoclonal antibody (CL647-66194, Clone:5E3A6) (red), or 0.4 ug CoraLite® Plus 647 Mouse IgG2a Isotype Control (C1.18.4) (CL647-65208, Clone: C1.18.4) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).