For Research Use Only

CoraLite® Plus 647-conjugated Phospho-YAP1 (Ser127) Recombinant antibody




Catalog Number:CL647-80694-2

Basic Information	Catalog Number: CL647-80694-2	GenBank Accession Number: BC038235	Purification Method: Protein A purification	
	Concentration: 1000 µg/ml	GenelD (NCBI): 10413	CloneNo.: 240779G4	
	Source: Rabbit	UNIPROT ID: P46937 Full Name: Yes-associated protein 1, 65kDa Calculated MW: 504 aa, 54 kDa	Excitation/Emission maxima wavelengths: 654 nm / 674 nm	
	lsotype: IgG			
		Observed MW: 70-78 kDa		
Applications	Tested Applications: FC (Intra)			
	Species Specificity: human			
Background Informati	of the Hippo pathway, and the translational modifications cytoplasmic retention of YA degradation that is depended phosphorylation of YAP at S	The transcriptional factors Yes-associated protein 1 (YAP1) and PDZ-binding motif (TAZ) act as downstream effectors of the Hippo pathway, and their subcellular location and transcriptional activities are affected by multiple post- translational modifications (PTMs). After phosphorylation, YAP1/TAZ binds to the 14-3-3 protein, which induces cytoplasmic retention of YAP1/TAZ. Moreover, phosphorylated YAP1/TAZ is likely to undergo ubiquitination degradation that is dependent on the β -transducin repeat-containing E3 ubiquitin protein ligase complex. The phosphorylation of YAP1 Ser127 promotes its cytoplasmic retention, whereas phosphorylation at Ser397 induces degradation. (PMID: 29682330, PMID: 26039999)		
		Storage: Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer: PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3. Aliquoting is unnecessary for -20°C storage		

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1X10^6 HepG2 cells untreated (dashed lines) or treated with Calyculin A which intracellularly stained with 0.25 ug Coralite® Plus 647 Phospho-Yap1 (Ser127) Recombinant Antibody (CL647-80694-2, Clone:240779G4)(red), or 0.25 ug Coralite® Plus 647 Rabbit IgG Isotype Control RecAb (CL647-98136, Clone: 240953C9) (blue). Cells were fixed with 4% PFA and permeabilized with 90% MeOH.