

For Research Use Only

# CoraLite® Plus 750-conjugated HDAC3 Recombinant antibody



Catalog Number: CL750-81211

## Basic Information

<b>Catalog Number:</b> CL750-81211	<b>GenBank Accession Number:</b> BC000614	<b>Purification Method:</b> Protein A purification
<b>Size:</b> 1000 µg/ml	<b>GeneID (NCBI):</b> 8841	<b>CloneNo.:</b> 2N1
<b>Source:</b> Rabbit	<b>UNIPROT ID:</b> O15379	<b>Recommended Dilutions:</b> WB 1:500-1:2000
<b>Isotype:</b> IgG	<b>Full Name:</b> histone deacetylase 3	<b>Excitation/Emission maxima wavelengths:</b> 755 nm / 780 nm
<b>Immunogen Catalog Number:</b> AG0396	<b>Calculated MW:</b> 49 kDa	
	<b>Observed MW:</b> 49 kDa	

## Applications

<b>Tested Applications:</b> WB	<b>Positive Controls:</b> WB : HeLa cells, Jurkat cells
<b>Species Specificity:</b> human, mouse, rat	

## Background Information

Histone acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. Histone deacetylase (HDAC) and histone acetyltransferase (HAT) are enzymes that regulate transcription by selectively deacetylating or acetylating the (-amino groups of lysines located near the amino termini of core histone proteins. At least 4 classes of HDAC were identified. HDAC3 is a class I HDAC. HDAC3 has histone deacetylase activity and may participate in the regulation of transcription through its binding with the zinc-finger transcription factor YY1. HDAC3 can also down-regulate p53 function and thus modulate cell growth and apoptosis. The gene encoding HDAC3 is regarded as a potential tumor suppressor gene.

## Storage

**Storage:**  
Store at -20°C. Avoid exposure to light. Stable for one year after shipment.  
**Storage Buffer:**  
PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.  
**Aliquoting is unnecessary for -20°C storage**

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)

W: [ptgcn.com](http://ptgcn.com)

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with CL750-81211 (HDAC3 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.