For Research Use Only

CoraLite® Plus 750-conjugated spike protein Recombinant antibody

Proteintech

Antibodies | ELISA kits | Proteins

www.ptglab.com

Catalog Number: CL750-82734-4

Basic Information

Catalog Number:

CL750-82734-4

Size: 1000 µg/ml Source:

Isotype:

Immunogen Catalog Number:

AG30684

Rabbit

GenBank Accession Number:

NC_045512 GeneID (NCBI): 43740568 Full Name:

SARS-CoV-2 Spike Protein

Calculated MW: 141 kDa Purification Method: Protein A purification

CloneNo.: 4l18

Excitation/Emission maxima

wavelengths: 755 nm / 780 nm

Applications

Tested Applications:

Species Specificity:

virus

Background Information

Coronaviruses (CoVs) infect human and animals and cause varieties of diseases, including respiratory, enteric, renal, and neurological diseases. CoV uses its spike protein to recognize ACE2 as its receptors and mediate membrane fusion and virus entry into host cells(PMID: 32221306). Each monomer of trimeric S protein is about 180 kDa, and contains two subunits, S1 and S2,S1 recognizes and binds to host receptors, and subsequent conformational changes in S2 facilitate fusion between the viral envelope and the host cell membrane(PMID: 19198616). Although the amino acid sequences of the S-glycoprotein were found to be different between the various HCoV, the structures showed high similarity, but the best 3D structural overlap shared by SARS-CoV and SARS-CoV-2, consistent with the shared ACE2 predicted receptor (PMID: 32522207). The spike protein of CoVs can be a target for vaccine and therapeutic development (PMID: 19198616). 28901-1-AP is specific for spike protein of SARS-COV-2, that antigen region is 428-506aa.

Storage

Storage:


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with CL750-82734-4 (spike protein antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.