For Research Use Only

## CoraLite® Plus 750-conjugated PTPN18 Recombinant antibody

Catalog Number: CL750-83024



**Purification Method:** 

Protein A purification

Recommended Dilutions:

Excitation/Emission maxima

CloneNo.:

230154G8

WB 1:500-1:2000

wavelengths: 755 nm / 780 nm

**Basic Information** 

Catalog Number: CL750-83024

Size: 1000 ug/ml Source: Rabbit Isotype:

Immunogen Catalog Number:

AG34048

**Tested Applications:** WB, FC (Intra) Species Specificity:

GenBank Accession Number:

BC052800 GeneID (NCBI): 26469 **UNIPROT ID:** Q99952

Full Name: protein tyrosine phosphatase, nonreceptor type 18 (brain-derived)

Calculated MW: 460 aa, 50 kDa Observed MW: 48 kDa

Positive Controls:

WB: HepG2 cells, MCF-7 cells, T-47D cells

## **Background Information**

PTPN18 (also called PTP-HSCF or BDP1), a member of the protein tyrosine phosphatase superfamily, has been  $reported \ to \ be \ a \ tumor \ suppressor \ in \ breast \ cancer \ by \ dephosphory lating \ HER2 \ and \ suppressing \ cell \ proliferation \ and \ suppressing \ cell \ proliferation \ and \ suppression \ cell \ proliferation \ and \ suppression \ s$ invasion. In addition to the well-established role of PTPN18 in breast cancer, our previous study found that PTPN18 and CSK cooperate to inhibit the events triggered by SRC kinases. (PMID: 35982039)

Storage


**Applications** 

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

## **Selected Validation Data**



Various lysates were subjected to SDS PAGE followed by western blot with CL750-83024 (PTPN18 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



1x10^6 MCF-7 cells were intracellularly stained with 0.25 ug CoraLite® Plus 750-conjugated PTPN18 Recombinant antibody (CL750-83024, Clone:230154G8)(red), or 0.25 ug Rabbit 1gG Isotype Control Recombinant Antibody (98136-1-RR, Clone: 240953C9) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).