For Research Use Only AGI-6780



www.ptgcn.com

Catalog Number: CM04694

产品信息

Catalog Number: CM04694


CAS号: 1432660-47-3

分子式: C₂₁H₁₈F₃N₃O₃S₂

主要靶点: Isocitrate Dehydrogenase (IDH)|Dehydrogenase

主要通路: 代谢 分子量: 481.51 溶解度:

DMSO:89 mg/mL (184.8 mM),Ethanol:89 mg/mL (184.8 mM),H2O:<1 mg/mL



靶点活性

IDH2 R140Q mutant:23 nM

体内活性

AGI-6780逆转TF-1细胞中IDH2/R140Q诱导的分化阻滞,并诱导原发性人类IDH2/R140Q AML患者样品中的急剧分化。AGI-6780有效降低异位过表达IDH2/R140Q的细胞系中的(R)-2-羟基戊二酸(2 hG)水平,EC 50为20 nM,与其他脱氢酶相比,具有优异的选择性。

细胞实验

AGI-6780 is dissolved in DMSO and stored, and then diluted with appropriate media before use[1]. Cells are sorted from fresh or frozen bone marrow aspirates and blood samples after labelling with PE-CD34, APC-CD38, PE-CD14, FITC-CD3 (clone HIT3a) and PECy7-CD19 (clone SJ25C1) antibodies using a MoFlow cell sorter. Unfractionated nucleated blood or bone marrow cells are plated in Methocult H4434 methylcellulose medium at 104 cells/dish, in duplicate dishes per condition. AGI-6780 (5 mM) is directly added to the medium. Dishes are incubated in a humidified incubator at 37°C and colonies containing at least 30 cells are counted after 13 days[1].

描试

AGI-6780 is a potent and selective inhibitor of IDH2 R140Q mutant.

储存

Powder: -20°C for 3 years | In solvent: -80°C for 2 years