For Research Use Only

PerCP-Cyanine 5.5 Anti-Mouse CD8a (53-6.7)



Catalog Number: CPY5-65069

Basic Information

Catalog Number:

GenBank Accession Number:

Purification Method: Affinity purification

CPY5-65069

GeneID (NCBI):

BC030679

CloneNo.:

12525

100ug, 100 $\,\mu$ g/ml Source:

UNIPROT ID:

Excitation/Emission maxima

Isotype:

P01731 Full Name: wavelengths: 482 nm / 677 nm

IgG2a, kappa

CD8 antigen, alpha chain

Applications

Tested Applications:

Species Specificity:

Mouse

Background Information

CD8 is a transmembrane glycoprotein composed of two disulfide-linked chains. It can be present as a homodimer of CD8 α or as a heterodimer of CD8 α and CD8 β (PMID: 3264320; 8253791). CD8 is found on most thymocytes. The majority of class I-restricted T cells express mostly the CD8 α β heterodimer while CD8 α α homodimers alone have been found on some gut intraepithelial T cells , on some T cell receptor (TCR) γ δ T cells and on NK cells (PMID: 2111591; 1831127; 8420975). CD8 acts as a co-receptor that binds to MHC class-I and participates in cytotoxic T cell activation (PMID: 8499079). During T cell development, CD8 is required for positive selection of CD4-/CD8+T cells (PMID: 1968084).


Storage

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:

PBS with 0.09% sodium azide and 0.5% BSA.

Selected Validation Data



1x10^6 mouse splenocytes were surface stained with 0.25 ug PerCP-Cyanine5.5 Anti-Mouse CD8a (53-6.7) (CPY5-65069, Clone:53-6.7) or unstained. Cells were then stained with 0.25 ug CoraLite® Plus 647 Anti-Mouse CD4 (GK1.5) (CL647-65104, Clone: GK1.5). Cells were not fixed.