

For Research Use Only

Recombinant Human CD40L/CD154 protein (His Tag)



Catalog Number: Eg0054

Basic Information

ED50:
17-68 ng/mL

GeneID:
959

Species:
Human

Accession:
P29965

Purity:
>95 %, SDS-PAGE

Technical Specifications

Purity:
>95 %, SDS-PAGE

Endotoxin Level:
<1.0 EU/ μ g protein, LAL method

Source:
HEK293-derived Human CD40L protein Met113-Leu261 (Accession# P29965) with a His tag at the N-terminus.

Predicted Molecular Mass:
17 kDa

SDS-PAGE:
17-19 kDa, reducing (R) conditions

Formulation:
Lyophilized from sterile PBS, pH 7.4. Normally 5% trehalose and 5% mannitol are added as protectants before lyophilization.

Biological Activity

Immobilized Human CD40L (His tag) at 2 μ g/mL (100 μ L/well) can bind Human CD40 (Myc tag, His tag) with a linear range of 17-68 ng/mL.

Storage and Shipping

Storage:
It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

- Until expiry date, -20°C to -80°C as lyophilized proteins.
- 3 months, -20°C to -80°C under sterile conditions after reconstitution.

Shipping:
The product is shipped at ambient temperature. Upon receipt, store it immediately at the recommended temperature.

Reconstitution

Briefly centrifuge the tube before opening. Reconstitute at 0.1-0.5 mg/mL in sterile water.

Background

The CD40 ligand (CD40L), also known as TRAP or CD154, is a member of the TNF superfamily of ligands. CD40L is primarily expressed on activated CD4⁺ T cells and on a small proportion of CD8⁺ T cells and platelets. It binds to CD40 on antigen-presenting cells (APC), which leads to many effects depending on the target cell type. It has been suggested that CD40/CD40L interactions regulate oxidative stress and affect various signaling pathways in both the immunological and the cardiovascular systems. The CD40/CD40L system is also involved in tumorigenesis. Its expression is tightly regulated, and abnormal levels of CD40L are associated with the pathogenesis of atheromatous plaque destabilization and thrombotic events. Multiple mutations in CD40LG gene have been identified that are associated with hyper-IgM immunodeficiency syndrome type 1.


References

1. Elgueta R. et al. (2009). Immunol Rev. 229(1):152-172.
2. Rizvi M. et al. (2008). Trends Mol Med. 14(12):530-538.
3. Michel N A. et al. (2017). Front Cardiovasc Med. 4:40.


Synonyms

CD154, CD40 Ligand, CD40L, CD40LG, gp39, HIGM1, IGM, IMD3, T-BAM, TNFSF5, TRAP

Selected Validation Data



Purity of Recombinant Human CD40L was determined by SDS-PAGE. The protein was resolved in an SDS-PAGE in reducing (R) and non-reducing (NR) conditions and stained using Coomassie blue.



Immobilized Human CD40L (His tag) at 2 μ g/mL (100 μ L/well) can bind Human CD40 (Myc tag, His tag) with a linear range of 17-68 ng/mL.

For technical support and original validation data for this product please contact

T: 027-87531629

E: Proteintech-CN@ptglab.com

W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.