

For Research Use Only

# Recombinant Mouse IL-2 protein (hFc Tag)



Catalog Number: Eg0456

## Basic Information

**ED50:**  
5-20 ng/mL

**GeneID:**  
16183

**Species:**  
Mouse

**Accession:**  
P04351

**Purity:**  
>90 %, SDS-PAGE

## Technical Specifications

**Purity:**  
>90 %, SDS-PAGE

**Endotoxin Level:**  
<1.0 EU/  $\mu$ g protein, LAL method

**Source:**  
HEK293-derived Mouse IL-2 protein Met+Ala21-Gln169 (Accession# P04351) with a human IgG1 Fc tag at the N-terminus.

**Predicted Molecular Mass:**  
43.5 kDa

**SDS-PAGE:**  
45-50 kDa, reducing (R) conditions

**Formulation:**  
Lyophilized from sterile PBS, pH 7.4. Normally 5% trehalose and 5% mannitol are added as protectants before lyophilization.

## Biological Activity

Measured in a cell proliferation assay using CTLL-2 mouse cytotoxic T cells. The ED50 for this effect is 5-20 ng/mL.

## Storage and Shipping

**Storage:**  
It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

- Until expiry date, -20°C to -80°C as lyophilized proteins.
- 3 months, -20°C to -80°C under sterile conditions after reconstitution.

**Shipping:**  
The product is shipped at ambient temperature. Upon receipt, store it immediately at the recommended temperature.

## Reconstitution

Briefly centrifuge the tube before opening. Reconstitute at 0.1-0.5 mg/mL in sterile water.

## Background

IL-2 (Interleukin-2) is an interleukin, a type of cytokine signaling molecule in the immune system. It regulates white blood cells (leukocytes, often lymphocytes) that are responsible for immunity. The major sources of IL-2 are activated CD4+ T cells and activated CD8+ T cells. IL-2 mediates its effects by binding to IL-2 receptors which are expressed by lymphocytes. IL-2 signal can be transduced by 3 different signaling pathways: JAK/STAT, PI3K/Akt/mTOR and MAPK/ERK pathway. IL-2 plays a role in regulating the adaptive immune system by controlling the survival and proliferation of regulatory T-cells, which are required for the maintenance of immune tolerance. It also promotes proliferation of activated B cells and subsequently immunoglobulin production.


## References

- 1.M C Mingari. et al. (1984). Nature. 312(5995): 641-3.
- 2.Thomas R Malek. et al. (2010). Immunity. 33(2): 153-65.
- 3.Wei Liao. et al. (2011). Curr Opin Immunol. 23(5): 598-604.
- 4.Natalia Arenas Ramirez. et al. (2015). Trends Immunol. 36(12): 763-777.


## Synonyms

IL-2, Interleukin 2, Lymphokine, TCGF

## Selected Validation Data



Purity of Recombinant Mouse IL-2 was determined by SDS-PAGE. The protein was resolved in an SDS-PAGE in reducing (R) and non-reducing (NR) conditions and stained using Coomassie blue.



Measured in a cell proliferation assay using CTLL-2 mouse cytotoxic T cells. The ED50 for this effect is 5-20 ng/mL.

For technical support and original validation data for this product please contact

T: 027-87531629

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)

W: [ptgcn.com](http://ptgcn.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.