

For Research Use Only

# Recombinant Human TNFR1/CD120a protein (His Tag)



Catalog Number: Eg0531

## Basic Information

**ED50:**  
0.8-3.2 ng/mL

**GeneID:**  
7132

**Species:**  
Human

**Accession:**  
P19438-1

**Purity:**  
>95 %, SDS-PAGE

## Technical Specifications

**Purity:**  
>95 %, SDS-PAGE

**Endotoxin Level:**  
<1.0 EU/  $\mu$ g protein, LAL method

**Source:**  
HEK293-derived Human TNFR1 protein Ile22-Thr211 (Accession# P19438-1) with a His tag at the C-terminus.

**Predicted Molecular Mass:**  
24 kDa

**SDS-PAGE:**  
32-48 kDa

**Formulation:**  
Lyophilized from sterile PBS, pH 7.4. Normally 5% trehalose and 5% mannitol are added as protectants before lyophilization.

## Biological Activity

Immobilized Human TNFR1 (His tag) at 0.2  $\mu$ g/mL (100  $\mu$ L/well) can bind Human TNF alpha (GST tag) with a linear range of 0.8-3.2 ng/mL.

## Storage and Shipping

**Storage:**  
It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

- Until expiry date, -20°C to -80°C as lyophilized proteins.
- 3 months, -20°C to -80°C under sterile conditions after reconstitution.

**Shipping:**  
The product is shipped at ambient temperature. Upon receipt, store it immediately at the recommended temperature.

## Reconstitution

Briefly centrifuge the tube before opening. Reconstitute at 0.1-0.5 mg/mL in sterile water.

## Background

Tumor necrosis factor (TNF) is a multifunctional cytokine that plays a key role in regulating inflammation, immune functions, host defense, and apoptosis. TNF signals through two distinct cell surface receptors, TNFR1 (TNFRSF1A, CD120a) and TNFR2 (TNFRSF1B, CD120b). TNFR1, which contains a death domain (DD) within its intracytoplasmic region, is thought to be the key receptor for TNF signaling. This receptor can be released to the extracellular space by two mechanisms, ectodomain shedding and constitutive release of full-length 55 kDa TNFR1. Soluble TNFR1 (sTNFR1) could function as TNF-binding protein, inhibiting TNF bioactivity.


## References

1. Islam A, et al. (2006). J Biol Chem. 281(10):6860-73.
2. Aggarwal BB, et al. (2012). Blood. 119(3):651-65.
3. Aderka D, et al. (2012). J Exp Med. 175(2):323-9.


## Synonyms

CD120a, FPF, p55, p55-R, p60, TBP1, TNF RI, TNFAR, TNF-R, TNF-R1, TNFR55, TNF-R55, TNFR60, TNF-R-I, TNFRSF1A

## Selected Validation Data



Purity of Recombinant Human TNFR1 was determined by SDS-PAGE. The protein was resolved in an SDS-PAGE in reducing (R) conditions and stained using Coomassie blue.



Immobilized Human TNFR1 (His tag) at 0.2  $\mu$ g/mL (100  $\mu$ L/well) can bind Human TNF alpha (GST tag) with a linear range of 0.8-3.2 ng/mL.

For technical support and original validation data for this product please contact

T: 027-87531629

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)

W: [ptgcn.com](http://ptgcn.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.