

For Research Use Only

Recombinant Human CD28 protein (hFc Tag)



Catalog Number: Eg0969

Basic Information

ED50:
6-24 ng/mL

GenelD:
940

Species:
Human

Accession:
P10747-1

Purity:
>90 %, SDS-PAGE

Technical Specifications

Purity:
>90 %, SDS-PAGE

Endotoxin Level:
<1.0 EU/ μ g protein, LAL method

Source:
HEK293-derived Human CD28 protein Asn19-Pro152 (Accession# P10747-1) with a human IgG1 Fc tag at the C-terminus.

Predicted Molecular Mass:
41.1 kDa

SDS-PAGE:
52-65 kDa, reducing (R) conditions

Formulation:
Lyophilized from sterile PBS, pH 7.4. Normally 5% trehalose and 5% mannitol are added as protectants before lyophilization.

Biological Activity

Immobilized Human CD80 (Myc tag, His tag) at 2 μ g/mL (100 μ L/well) can bind Human CD28 (hFc tag) with a linear range of 6-24 ng/mL.

Storage and Shipping

Storage:
It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

- Until expiry date, -20°C to -80°C as lyophilized proteins.
- 3 months, -20°C to -80°C under sterile conditions after reconstitution.

Shipping:
The product is shipped at ambient temperature. Upon receipt, store it immediately at the recommended temperature.

Reconstitution

Briefly centrifuge the tube before opening. Reconstitute at 0.1-0.5 mg/mL in sterile water.

Background

Cluster of Differentiation 28 (CD28, also known as Tp44) is one of the proteins constitutively expressed on the surface of 80% of human T cells. CD28 is a highly glycosylated membrane protein, formed by two homodimer disulfide-linked peptides, which shares homology with the immunoglobulin superfamily. CD28 signals play a key role in many T cell processes including cytoskeletal remodeling, production of cytokines, survival, and differentiation. CD28 interactions with the B7 family of costimulatory ligands are essential for initiating antigen-specific T cell responses, upregulating cytokine expression and promoting T cell expansion and differentiation.


References

- 1.Hansen JA. et al. (1980). Immunogenetics. 10(1-4): 247-260.
- 2.Aruffo A. et al. (1987). Proc Natl Acad Sci U S A. 84(23): 8573-8577.
- 3.Jonathan H Esensten. et al. (2016). Immunity. 44(5):973-988.
- 4.DJ Lenschow. et al. (1996). Annu Rev Immunol. 14:233-258.


Synonyms

CD28, Tp44

Selected Validation Data



Purity of Recombinant Human CD28 was determined by SDS-PAGE. The protein was resolved in an SDS-PAGE in reducing (R) and non-reducing (NR) conditions and stained using Coomassie blue.



Immobilized Human CD80 (Myc tag, His tag) at $2 \mu\text{g/mL}$ ($100 \mu\text{L/well}$) can bind Human CD28 (hFc tag) with a linear range of 6-24 ng/mL.

For technical support and original validation data for this product please contact

T: 027-87531629

E: Proteintech-CN@ptglab.com

W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.