


Recombinant Human BST2 protein (rFc Tag)

Catalog Number: Eg1961

Basic Information	Species: Human	Purity: >90 %, SDS-PAGE	Tag: rFc Tag
Technical Specifications	Purity: >90 %, SDS-PAGE Endotoxin Level: <0.1 EU/ µg protein, LAL method Source: HEK293-derived Human BST2 protein Asn49-Ser161 (Accession# Q10589-1) with a rabbit IgG Fc tag at the N-terminus. GeneID: 684 Accession: Q10589-1 Predicted Molecular Mass: 39.8 kDa SDS-PAGE: 40-50 kDa, reducing (R) conditions Formulation: Lyophilized from 0.22 µm filtered solution in PBS, pH 7.4. Normally 5% trehalose and 5% mannitol are added as protectants before lyophilization.		
Biological Activity	Not tested		
Storage and Shipping	Storage: It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles. • Until expiry date, -20°C to -80°C as lyophilized proteins. • 3 months, -20°C to -80°C under sterile conditions after reconstitution. Shipping: The product is shipped at ambient temperature. Upon receipt, store it immediately at the recommended temperature.		
Reconstitution	Briefly centrifuge the tube before opening. Reconstitute at 0.1-0.5 mg/mL in sterile water.		
Background	BST2, also named as CD317 and Tetherin, belongs to the tetherin family. It may be involved in the sorting of secreted proteins and it is involved in pre-B-cell growth. BST2 is an antiretroviral defense protein, that blocks retrovirus release from the cell surface. Depleted upon HIV-1 infection by viral VPU protein through 20S proteasome degradation. Depleted upon infection by human Kaposi's sarcoma-associated herpesvirus (KSHV) through ubiquitination and subsequent degradation. BST2 may play a role in B-cell activation in rheumatoid arthritis. It is recently identified interferon-induced cellular proteins that restrict infections by retroviruses and filoviruses and of influenza virus and flaviviruses, respectively. BST2 is a plasma membrane protein, tetherin inhibits virion particle release from infected cells. BST2 is effective against retroviruses and flaviviruses whilst IFITMs disrupt influenza and flavivirus infection.		
References	1. Miyagi E. et al. (2009). Proc Natl Acad Sci U S A. 106(8):2868-2873. 2. Skasko M. et al. (2011). Virology. 411(1):65-77. 3. Andrew AJ. et al. (2009). Retrovirology. 6:80. 4. Tanwattana N. et al. (2023). PLoS One. 18(11):e0292833.		
Synonyms	CD317, Bone marrow stromal antigen 2, BST 2, BST-2, CD 317		

Selected Validation Data



Purity of Recombinant Human BST2 was determined by SDS-PAGE. The protein was resolved in an SDS-PAGE in reducing (R) conditions and stained using Coomassie blue.

For technical support and original validation data for this product please contact

T: 027-87531629

E: Proteintech-CN@ptglab.com

W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.