For Research Use Only

Recombinant human GPR56 protein (rFc Tag)



www.ptgcn.com

Catalog Number: Eg2478

Basic Information

ED50:

Species:

Purity: >90 %, SDS-PAGE

GeneID: 9289

Accession: Q9Y653-1

Technical Specifications

Purity: >90 %, SDS-PAGE

Endotoxin Level: <1.0 EU/ μ g protein, LAL method

HEK293-derived Human GPR56 protein Arg26-Asn171 (Accession# Q9Y653-1) with a rabbit IgG Fc tag at the C-

Predicted Molecular Mass:

42.7 kDa

SDS-PAGE:

45-55 kDa, reducing (R) condition

Lyophilized from sterile PBS, pH 7.4. Normally 5% trehalose and 5% mannitol are added as protectants before

lyophilization.

Biological Activity

Not tested

Storage and Shipping

It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Until expiry date, -20°C to -80°C as lyophilized proteins.

3 months, -20℃ to -80℃ under sterile conditions after reconstitution.

The product is shipped at ambient temperature. Upon receipt, store it immediately at the recommended

temperature.

Reconstitution

Briefly centrifuge the tube before opening. Reconstitute at 0.1-0.5 mg/mL in sterile water.

Background

GPR56 belongs to the adhesion family of G protein-coupled receptors (GPRs), is a receptor involved in cell adhesion and probably in cell-cell interactions, and mediates cell matrix adhesion in developing neurons and hematopoietic stem cells. GPR56 binding to the COL3A1 ligand inhibits neuronal migration and activates the RhoA pathway by coupling to GNA13 and possibly GNA12. GPR56 Plays a critical role in cancer progression by inhibiting VEGFA production thereby inhibiting angiogenesis through a signaling pathway mediated by PRKCA.


References

1. Rong Luo, et al. (2012) PLoS One. 7(1):e29818. 2. Lei Xu, et al. (2006) Proc Natl Acad Sci U S A. 103(24):9023-8. 3. Liquan Yang, et al. (2011) Cancer Res. 71(16):5558-68.

Synonyms

GPR56,ADGRG1,ADGRG1 CT,ADGRG1 C-terminal fragment,

Selected Validation Data



Purity of Recombinant Human GPR56 was determined by SDS-PAGE. The protein was resolved in an SDS-PAGE in reducing (R) and non-reducing (NR) conditions and stained using Coomassie blue.