For Research Use Only

FITC Plus-conjugated HIF-1 alpha Polyclonal antibody

proteintech®
Antibodies | ELISA kits | Proteins
www.ptglab.com

Catalog Number:FITC-20960 Featured Product

Basic Information

Catalog Number:
FITC-20960

Size:
1000 μg/ml

Source:
Rabbit

Isotype:
IgG

Immunogen Catalog Number:
AG15198

GenBank Accession Number:
BC012527

GeneID (NCBI):
3091

UNIPROT ID:
Q16665

Full Name:
hypoxia inducible factor 1, alpha subunit (basic helix-loop-helix transcription factor)

Calculated MW:
826 aa, 93 kDa

Observed MW:
120 kDa

Purification Method:
Antigen affinity purification

Recommended Dilutions:
IF/ICC 1:50-1:500

Excitation/Emission maxima wavelengths:
495 nm / 524 nm

Applications

Tested Applications:
IF/ICC

Species Specificity:
human

Positive Controls:

IF/ICC : Cobalt Chloride treated HeLa cells,

Background Information

HIF1a, the major regulator of the cellular responses to hypoxia, consists of an oxygen-sensitive subunit, HIF1 alpha (HIF1A), and an oxygen-insensitive subunit, HIF1 beta (arylhydrocarbon receptor nuclear transporter [ARNT]). Under normal oxygen conditions, HIF1a is continuously produced and destroyed, in a process involving hydroxylation, interaction with von Hippel-Lindau (VHL) protein, polyubiquitylation and subsequent proteasomal degradation. Under hypoxic conditions, hydroxylation is impaired and HIF1a is stabilized. HIF1a localizes in cytoplasm in normoxia, but it can translocate into nuclear in response to hypoxia. The calculated molecular weight of HIF1a is 93 kDa, but the modified protein HIF1a is about 110-120kDa (PMID: 11698256, .PMID: 7539918). .

Storage

Storage:
Store at -20°C. Avoid exposure to light.
Storage Buffer:
PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:
T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com

Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed Cobalt Chloride treated HeLa cells using FITC Plus HIF-1 alpha antibody (FITC-20960) at dilution of 1:50.