For Research Use Only

FITC Plus-conjugated ATF4 Monoclonal antibody

Catalog Number:FITC-60035



Basic Information	Catalog Number: FITC-60035	GenBank Accession Number: BC022088	Purification Method: Protein G purification
	Size: 1000 µg/ml	GenelD (NCBI): 468	CloneNo.: 3E4C5
	Source: Mouse	UNIPROT ID:Excitation/Emission maximaP18848wavelengths:	
	Isotype: IgG1 Immunogen Catalog Number: AG1279	Full Name: activating transcription factor 4 (tax- responsive enhancer element B67)	hancer element B67) V:
		Calculated MW: 39 kDa	
		Observed MW: 45-50 kDa	
Applications	Tested Applications: FC (Intra)		
	Species Specificity: human, mouse, rat		
Background Information	ATF4 is a transcription factor, that accumulates predominantly in osteoblasts, where it regulates terminal osteoblast differentiation and bone formation. As a basic leucine-zipper (bZip) transcription factor, ATF4 can regulate amino acid metabolism, cellular redox state, and anti-stress responses. It also regulates age-related and diet-induced obesity and glucose homeostasis in mammals, and has conserved metabolic functions in flies. Due to its location at		

differentiation and bone formation. As a basic leucine-zipper (bZip) transcription factor, ATF4 can regulate amino acid metabolism, cellular redox state, and anti-stress responses. It also regulates age-related and diet-induced obesity and glucose homeostasis in mammals, and has conserved metabolic functions in flies. Due to its location at chromosome 22q13, a region linked to schizophrenia, ATF4 is considered as a positional candidate gene for schizophrenia. Otherwise, since ATF4 is induced by tumour microenvironmental factors, and regulates processes relevant to cancer progression, it might serve as a potential therapeutic target in cancer. This antibody is a mouse monoclonal antibody raised against full length human ATF4 antigen. The antibody recognizes the 45-50 kDa ATF4 protein and its phosphorylated forms (50 kDa).

Storage


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer: PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3. Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1X10⁶ HepG2 cells were intracellularly stained with 0.4 ug FITC Plus Anti-Human ATF4 (FITC-60035, Clone:3E4C5) (red), or 0.4 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).