## For Research Use Only

## FITC Plus Anti-Mouse CD80 (B7-1) (16-10A1)




Catalog Number:FITC-65076

**8** Publications

Basic Information	Catalog Number: GenBank Accession Number: FITC-65076 BC131959		cession Number:	Purification Method: Affinity purification	
	<mark>Size:</mark> 100ug, 500 µg/ml	Genel D (NC) 12519	BI):	CloneNo.: 16-10A1	
	Source: Armenian Hamster	UNIPROT ID Q00609	:	Excitation/Emission maxima wavelengths:	
	Isotype: IgG	Full Name:495 nmCD80 antigen		495 nm / 524 nm	
Applications	Tested Applications: FC				
	Cited Applications: FC, IF				
	Species Specificity: Mouse				
	Cited Species: mouse				
	CD80 (also known as B7-1) is a type I membrane protein that is a member of the immunoglobulin superfamily, wi an extracellular immunoglobulin constant-like domain and a variable-like domain required for receptor binding. is expressed on antigen-presenting cells (APCs), including B cells, dendritic cells, monocytes, and macrophages. CD80 is the receptor for the proteins CD28 and CTLA-4 found on the surface of T-cells. It is involved in the costimulatory signal essential for T-lymphocyte activation. T-cell proliferation and cytokine production is induced by the binding of CD28, binding to CTLA-4 has opposite effects and inhibits T-cell activation. CD80 also acts as a cellular attachment receptor for adenovirus subgroup B. (PMID: 7545666; 12015893; 16920215)				
Background Information	is expressed on antigen-pre CD80 is the receptor for the costimulatory signal essent by the binding of CD28, bind	proteins CD28 and CTL ial for T-lymphocyte a ding to CTLA-4 has opp	ncluding B cells, dendri A-4 found on the surfac ctivation. T-cell prolife osite effects and inhibi	tic cells, monocytes, and macrophages. The of T-cells. It is involved in the ration and cytokine production is induce ts T-cell activation. CD80 also acts as a	
	is expressed on antigen-pre CD80 is the receptor for the costimulatory signal essent by the binding of CD28, bind	proteins CD28 and CTL ial for T-lymphocyte a ding to CTLA-4 has opp	ncluding B cells, dendri A-4 found on the surfac ctivation. T-cell prolife osite effects and inhibi	tic cells, monocytes, and macrophages. The of T-cells. It is involved in the ration and cytokine production is induce ts T-cell activation. CD80 also acts as a	
	is expressed on antigen-pre CD80 is the receptor for the costimulatory signal essent by the binding of CD28, bind cellular attachment recepto	proteins CD28 and CTL ial for T-lymphocyte a ding to CTLA-4 has opp r for adenovirus subgro	ncluding B cells, dendri A-4 found on the surfac ctivation. T-cell prolife osite effects and inhibi oup B. (PMID: 7545666;	tic cells, monocytes, and macrophages. The of T-cells. It is involved in the ration and cytokine production is induce ts T-cell activation. CD80 also acts as a 12015893; 16920215)	
	is expressed on antigen-pre CD80 is the receptor for the costimulatory signal essent by the binding of CD28, bin cellular attachment recepto	proteins CD28 and CTL ial for T-lymphocyte a ding to CTLA-4 has opp r for adenovirus subgro Pubmed ID	Actuding B cells, dendri A-4 found on the surfac ctivation. T-cell prolife osite effects and inhibi pup B. (PMID: 7545666; Journal	tic cells, monocytes, and macrophages. The of T-cells. It is involved in the ration and cytokine production is induce ts T-cell activation. CD80 also acts as a 12015893; 16920215) Application FC	
Background Information	is expressed on antigen-pre CD80 is the receptor for the costimulatory signal essent by the binding of CD28, bind cellular attachment recepto Author Xiaojin Chen	proteins CD28 and CTL ial for T-lymphocyte ad ding to CTLA-4 has opp r for adenovirus subgro Pubmed ID 36297436	Actuding B cells, dendri A-4 found on the surfac ctivation. T-cell prolife osite effects and inhibi pup B. (PMID: 7545666; Journal Pharmaceutics	tic cells, monocytes, and macrophages. The of T-cells. It is involved in the ration and cytokine production is induce ts T-cell activation. CD80 also acts as a 12015893; 16920215) Application FC	

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



1X10^6 Balb/c mouse peritoneal macrophages were surface stained with 0.5 ug FITC Anti-Mouse CD80 (B7-1) (FITC-65076, Clone:16-10A1) (red), or 0.5 ug Isotype Control. Cells were not fixed.