

For Research Use Only

# FITC Anti-Mouse CD16 / CD32 (2.4G2)



Catalog Number: FITC-65080

4 Publications

## Basic Information

Catalog Number:

FITC-65080

Size:

100ug, 0.5 mg/ml

Source:

Rat

Isotype:

IgG2b, kappa

GenBank Accession Number:

BC038070

GeneID (NCBI):

14130

UNIPROT ID:

P08101

Full Name:

Fc receptor, IgG, low affinity IIb

Purification Method:

Affinity purification

CloneNo.:

2.4G2

Excitation/Emission maxima wavelengths:

498 nm / 526 nm

## Applications

Tested Applications:

FC

Cited Applications:

FC

Species Specificity:

Mouse

Cited Species:

mouse

## Background Information

CD16 (Fc  $\gamma$  RIII) is a 50-70 kDa low affinity Fc receptor found on the surface of natural killer cells, neutrophil polymorphonuclear leukocytes, monocytes and macrophages. CD16 mediates antibody-dependent cellular cytotoxicity (ADCC) and other antibody-dependent responses, such as phagocytosis. CD32 (Fc  $\gamma$  RII) is a 40 kD transmembrane glycoprotein that binds to the Fc region of IgG with low affinity. CD32 is present on phagocytic cells such as macrophages and neutrophils, and is involved in the process of phagocytosis and clearing of immune complexes. The 2.4G2 antibody reacts with the extracellular domains of mouse CD16 (FcRIII) and CD32 (FcRII) (PMID: 90108; 8406898).

## Notable Publications

Author	Pubmed ID	Journal	Application
Qi Wang	34214019	Pharm Biol	FC
Yan Shi	38247194	Small	FC
Xinyi Qu	37445610	Int J Mol Sci	FC

## Storage

Storage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:

Phosphate based buffer with 0.09% sodium azide and 0.1% gelatin, pH 7.2.

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)

W: [ptgcn.com](http://ptgcn.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



1X10<sup>6</sup> C57BL/6 mouse splenocytes were surface stained with APC-Anti-Mouse CD3 $\epsilon$  (APC-65060, Clone: 145-2C11) and 0.25  $\mu$ g FITC-Rat IgG2b isotype control (left) or 0.25  $\mu$ g FITC Anti-Mouse CD16 / CD32 (FITC-65080, Clone: 2.4G2) (right). Cells were not fixed.