## For Research Use Only

## FITC Plus Anti-Human MICA/MICB (6D4)



Catalog Number:FITC-65161

1 Publications

**Basic Information** 

Catalog Number: FITC-65161

100tests, 5 µl/test

Source: Mouse

Size:

Isotype:

IgG2a, kappa

sequence A

Calculated MW: 383 aa, 43 kDa

GenBank Accession Number:

BC016929 GeneID (NCBI): 100507436 **UNIPROT ID:** Q29983 Full Name:

MHC class I polypeptide-related

**Purification Method:** 

The purified antibody is conjugated with fluorescein isothiocyanate (FITC) under optimum conditions. The conjugate is purified by sizeexclusion chromatography.

CloneNo.: 6D4

Excitation/Emission maxima wavelengths:

495 nm / 524 nm

**Applications** 

**Tested Applications:** 

Cited Applications:

Species Specificity:

**Background Information** 

Human MHC class I chain-related genes located within the HLA class I region of chromosome 6 encode MHC class I chain-related A and B (MICA and MICB) (PMID: 11429322). MICA and MICB are stress-inducible surface molecules that are not associated with \$2-microglobulin and do not present peptides (PMID: 9497295). They are expressed in intestinal epithelium and many epithelial tumors (PMID: 10359807). MICA and MICB are ligands for NKG2D which is an activating receptor that is expressed on most natural killer (NK) cells, CD8  $\alpha$   $\beta$  T cells, and  $\gamma$   $\delta$  T cells (PMID: 10426993; 11491531). The monoclonal antibody 6D4 recognizes MICA and MICB.

**Notable Publications** 


Author **Pubmed ID** Application Journal Tomoaki Naito 36660595 Biosci Microbiota Food Health FC

Storage


Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 0.1% sodium azide.

## Selected Validation Data



1X10^6 HeLa cells were surface stained with 5 ul FITC Plus Anti-Human MICA/MICB (FITC-65161, Clone:6D4) or Mouse IgG2a Isotype Control. Cells were not fixed. Live cells were gated.



1X10^6 HeLa cells were surface stained with 5 ul FITC Plus Anti-Human MICA/MICB (FITC-65161, Clone:6D4) or Mouse IgG2a Isotype Control. Cells were not fixed. Live cells were gated.