## For Research Use Only

## FITC Plus Anti-Rat CD8a (OX-8)

Catalog Number: FITC-65180



**Basic Information** 

Catalog Number:

GenBank Accession Number:

**Purification Method:** Purified by protein-A affinity

FITC-65180

GeneID (NCBI):

chromatography

Size:

24930

CloneNo.:

100ug, 500  $\,\mu$  g/ml Source:

UNIPROT ID: P07725

BC088126

ОХ-8

Mouse Isotype:

Full Name:

Excitation/Emission maxima wavelengths:

IgG1, kappa

CD8a molecule

495 nm / 524 nm

**Applications** 

**Tested Applications:** 

Species Specificity:

Rat

**Background Information** 

CD8 is a transmembrane glycoprotein composed of two disulfide-linked chains. It can be present as a homodimer of CD8  $\alpha$  or as a heterodimer of CD8  $\alpha$  and CD8  $\beta$  (PMID: 3264320; 8253791). CD8 is found on most thymocytes. The majority of class I-restricted T cells express mostly the CD8  $\alpha$   $\beta$  heterodimer while CD8  $\alpha$   $\alpha$  homodimers alone have been found on some gut intraepithelial T cells , on some T cell receptor (TCR)  $\gamma$   $\delta$  T cells and on NK cells (PMID: 2111591; 1831127; 8420975). CD8 acts as a co-receptor that binds to MHC class-I and participates in cytotoxic T cell activation (PMID: 8499079). During T cell development, CD8 is required for positive selection of CD4-/CD8+T cells (PMID: 1968084).


Storage

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:

PBS with 0.09% sodium azide.

## Selected Validation Data



1X10^6 Wistar rat splenocytes were surface stained with 0.5 ug FITC Plus Anti-Rat CD8a (FITC-65180, Clone: OX-8), or 0.5 ug Control Antibody. Cells were not fixed.