For Research Use Only

FITC Plus-conjugated CD64 Recombinant antibody

Catalog Number:FITC-65572



Purification Method:

Protein A purification

Excitation/Emission maxima

CloneNo.:

wavelengths: 495 nm / 524 nm

Basic Information

Catalog Number:

FITC-65572

Size:

100tests, 5 ul/test

Source: Mouse

Isotype:

lgG2a

374 aa, 43 kDa

GenBank Accession Number:

BC032634 GeneID (NCBI):

2209

ENSEMBL Gene ID: ENSG00000150337

Full Name:

Fc fragment of IgG, high affinity Ia,

receptor (CD64) Calculated MW:

Applications

Tested Applications:

Species Specificity:

human

Background Information


Fc γ receptor comprise a multigene family of integral membrane glycoproteins that exhibit complex activation or $inhibitory\ effects\ on\ cell\ functions\ after\ aggregation\ by\ complexed\ immunoglobulin\ G\ (IgG)\ (PMID:\ 17005690\).$ CD64, also known as Fc γ RIA, is a high-affinity receptor for the Fc region of IgG. It is expressed by monocytes/macrophages, activated neutrophils, dendritic cells, and early myeloid cells (PMID: 23293080; 19642859; 7680917). CD64 functions in both innate and adaptive immune responses.

Storage


Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer:

PBS with 0.09% sodium azide and 0.5% BSA.

Selected Validation Data



1x10^6 human PBMCs were surface stained with Coralite® Plus 647 Anti-Human CD14, and 5 ul FITC Plus Anti-Human CD64 (10.1) Mouse IgG2a Recombinant Antibody (FITC-65572, Clone:10.1) or FITC Plus Mouse IgG2a Isotype Control (C1.18.4) (FITC-65208, Clone: C1.18.4). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.



1x10^6 human PBMCs were surface stained with 5 ul FITC Plus Anti-Human CD64 (10.1) Mouse IgG2a Recombinant Antibody (FITC-65572, Clone:10.1) (red) or FITC Plus Mouse IgG2a Isotype Control (C1.18.4) (FITC-65208, Clone: C1.18.4) (blue). Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed.