For Research Use Only

HRP-conjugated GAPDH Monoclonal antibody



Catalog Number: HRP-60004

569 Publications

Basic Information

Catalog Number: HRP-60004

Size: 1000 µg/ml Source: Mouse

Isotype: IgG2b GenBank Accession Number: BC004109

BC004109
GeneID (NCBI):
2597
UNIPROT ID:
P04406
Full Name:

glyceraldehyde-3-phosphate dehydrogenase

Calculated MW: 36 kDa Observed MW: 36 kDa Purification Method:

Antigen affinity purification

CloneNo.: 1E6D9

Recommended Dilutions: WB 1:5000-1:50000

Applications

Tested Applications:

WB

Cited Applications: WB,IP,IHC,IF Species Specificity:

human, mouse, rat, zebrafish, plant

Cited Species:

human, Chicken, rat, Honey bees, mouse, rabbit,

hamster, pig, opossum

Positive Controls:

WB: HeLa cells, HepG2 cells, HEK-293 cells

Background Information

Glyceraldehyde-3-phosphate dehydrogenase (GAPDH) catalyzes the phosphorylation of glyceraldehyde-3-phosphate during glycolysis. GAPDH participates in nuclear events including transcription, binding RNA, RNA transportation, DNA replication, DNA repair and apoptosis. Being stably and constitutively expressed at high levels in most tissues and cells, GAPDH is considered a housekeeping protein. It is widely used as a control for RT-PCR and also loading control in electrophoresis and Western blotting. GAPDH is normally expressed in cellular cytoplasm or membrane, but can occasionally translocate to the nucleus after the addition of post-translational modifications such as S-nitrosylation. This antibody is raised against full length GAPDH of human origin. It can recognize the 36 kDa GAPDH protein in most cells/tissues. In addition, a band below 36 kDa can always be detected as the isoform or spliced product of GAPDH (PMID: 23885286, 23877755, 19368702). Please note that some physiological factors, such as hypoxia and diabetes, increase GAPDH expression in certain cell types.

Notable Publications

Author	Pubmed ID	Journal	Application
Dong Yu	36183792	Brain Res	WB
Jiawei Hao	36126167	Autophagy	WB
Victoria L. Messerschmidt	34651022	Front Cardiovasc Med	WB

Storage

Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926 E: Proteintech-CN@ptglab.com

W: ptgcn.co

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Western blot analysis of GAPDH in various tissues and cell lines using Proteintech antibody HRP-60004 at a dilution of 1:10000.