For Research Use Only

HRP-conjugated TDP-43 (human specific) Monoclonal antibody

www.ptglab.com

Catalog Number: HRP-60019

Featured Product

Basic Information

Catalog Number: GenBank Accession Number: BC001487

HRP-60019 GeneID (NCBI): Size: 1000 ug/ml 23435 **UNIPROT ID:** Source: Mouse Q13148 Full Name:

lgG1 TAR DNA binding protein

Calculated MW: Immunogen Catalog Number:

AG1231 43 kDa Observed MW: 43 kDa

Applications

Tested Applications:

WB

Species Specificity:

human

Isotype:

Purification Method:

Protein G purification

CloneNo.: 6H6E12

Positive Controls:

cells

Recommended Dilutions:

WB 1:5000-1:50000

WB: SH-SY5Y cells, HepG2 cells, Jurkat cells, K-562

Background Information

Transactivation response (TAR) DNA-binding protein of 43 kDa (also known as TARDBP or TDP-43) was first isolated as a transcriptional inactivator binding to the TAR DNA element of the HIV-1 virus. Neumann et al. (2006) found that a hyperphosphorylated, ubiquitinated, and cleaved form of TARDBP, known as pathologic TDP-43, is the major component of the tau-negative and ubiquitin-positive inclusions that characterize amyotrophic lateral sclerosis (ALS) and the most common pathological subtype of frontotemporal lobar degeneration (FTLD-U). Various forms of TDP-43 exist, including 18-35 kDa of cleaved C-terminal fragments, 45-50 kDa phospho-protein, 55 kDa glycosylated form, 75 kDa hyperphosphorylated form, and 90-300 kDa cross-linked form. (PMID: 17023659,19823856, 21666678, 22193176). 60019-2-lg is a mouse monoclonal antibody recognizing the cleavage $product\ of\ 20\text{-}30\ kDa\ in\ addition\ to\ the\ native\ and\ phosphory lated\ forms\ of\ TDP\text{-}43.\ Immunohistochemical\ analyses$ of TDP-43 using this antibody detect both normal diffuse nuclear staining and insoluble inclusions in pathologic tissues. Notably this antibody only recognizes human TDP-43 but not reacts with mouse or rat TDP-43.


Storage

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with HRP-60019 (TDP-43 (human specific) antibody) at dilution of 1:40000 incubated at room temperature for 1.5 hours.