## For Research Use Only

## HRP-conjugated OCT4 Monoclonal antibody



Catalog Number: HRP-60242

**Basic Information** 

Catalog Number: HRP-60242

Size:

1000 µg/ml

Source:

Mouse

Isotype:

IgG1 POU class 5 homeobox 1
Immunogen Catalog Number: Calculated MW:

AG1794 39 kDa

Observed MW: 50-52 kDa

BC020712

5460

GeneID (NCBI):

**UNIPROT ID:** 

Full Name:

Q01860

GenBank Accession Number:

Purification Method:

Protein G purification CloneNo.:

1C4B6

Recommended Dilutions: WB 1:5000-1:50000

**Applications** 

Tested Applications:

WB

Species Specificity: human, mouse

**Positive Controls** 

WB: HEK-293 cells, HT-29 cells, mouse brain tissue

## **Background Information**

Expression of the OCT4 transcription factor is associated with an undifferentiated phenotype of human and mouse embryonic stem cells and germ cells. OCT4 (Octamer-4) is a homeodomain transcription factor of the POU family (POU5F1) involved in regulation of pluripotency during normal development and is detectable in embryonic stem (ES) and germ cells. Its expression is associated with an undifferentiated phenotype and tumors. In fact, gene knockdown of OCT4 promotes differentiation, thereby demonstrating a role for this factor in human ES cell self-renewal. As such, it is frequently used as a marker for undifferentiated ES cells or iPS (induced pluripotent stem) cells. OCT4 is one of the four key transcription factors (OCT4, SOX2, c-MYC, and KLF4) used to reprogram mouse and human fibroblasts to a pluripotent state as well as generate patient-specific iPS cells. The calculated molecular weight of OCT4 is 38 kDa, but the modified OCT4 protein is about 50 kDa.

Storage


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer:


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



HEK-293 cells were subjected to SDS PAGE followed by western blot with HRP-60242 (OCT4 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with HRP-60242 (OCT4 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.