For Research Use Only

## HRP-conjugated GAPDH Recombinant antibody

Catalog Number: HRP-81640-5



**Basic Information** 

Catalog Number: HRP-81640-5

1000 µg/ml

Size:

BC004109

GeneID (NCBI):

GenBank Accession Number:

Source: UNIPROTID: Rabbit P04406

Isotype: Full Name:
IgG glyceraldehyde-3-phosphate

Immunogen Catalog Number: dehydrogenase

AG0766 Calculated MW: 36 kDa

Observed MW: 36 kDa Purification Method: Protein A purification

CloneNo.:

1H18

Recommended Dilutions: WB 1:2000-1:16000

**Applications** 

**Tested Applications:** 

WB

Species Specificity: human, mouse, rat

Positive Controls:

WB: HSC-T6 cells, rat brain tissue, mouse brain tissue

## **Background Information**

Glyceraldehyde-3-phosphate dehydrogenase (GAPDH) catalyzes the phosphorylation of glyceraldehyde-3-phosphate during glycolysis. GAPDH participates in nuclear events including transcription, binding RNA, RNA transportation, DNA replication, DNA repair and apoptosis. Being stably and constitutively expressed at high levels in most tissues and cells, GAPDH is considered a housekeeping protein. It is widely used as a control for RT-PCR and also loading control in electrophoresis and Western blotting. GAPDH is normally expressed in cellular cytoplasm or membrane, but can occasionally translocate to the nucleus after the addition of post-translational modifications such as S-nitrosylation. This antibody is raised against full length GAPDH of human origin. It can recognize the 36 kDa GAPDH protein in most cells/tissues. In addition, a band below 36 kDa can always be detected as the isoform or spliced product of GAPDH (PMID: 23885286, 23877755, 19368702). Please note that some physiological factors, such as hypoxia and diabetes, increase GAPDH expression in certain cell types.

Storage

Storage

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

torage Buffer

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with HRP-81640-5 (GAPDH antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.