

HumanKine[®] BMP-7 (Recombinant Human)



Animal Component-Free

Human cell expressed

Tag-Free

Endotoxin Free

Product Description

Animal-free Recombinant Human BMP-7 is expressed in human 293 cells as a disulfide linked homodimeric glycoprotein with an apparent molecular mass of 29 kDa. This cytokine is produced in a serum-free, chemically defined media. Recombinant Human BMP-7 is a homodimeric glycoprotein consisting of two 117 amino acid subunits, which correspond to amino acid residues 315 to 431 of the full-length BMP-7 precursor. The protein encoded by this gene is a member of the TGF-beta superfamily. Like other members of the Bone Morphogenetic Protein family of proteins, it plays a key role in the transformation of mesenchymal cells into bone and cartilage.


Alternative Names BMP 7, BMP7, BMP-7, bone morphogenetic protein 7, Eptoterminalfa, HZ1229, OP 1, OP1, Osteogenic protein 1

Source Human Embryonic Kidney cells (HEK293). HEK293-derived BMP-7 protein

Specifications


Test	Method	Specification
Activity	Dose-dependent induction of alkaline phosphatase production in the ATDC-5 cell line (mouse chondrogenic cell line)	50-275 ng/mL EC50
Molecular Mass	SDS-PAGE	20 kDa, homodimer, glycosylated
Purity	SDS-PAGE	>95%
Endotoxin	LAL	<1 EU/μg

Activity Data



Recombinant human BMP-7 (HZ-1229) stimulates dose-dependent induction of alkaline phosphatase production in the ATDC-5 mouse chondrogenic cell line. Alkaline phosphatase production was assessed using pNPP as a chromogenic substrate. ATDC-5 cells were treated with increasing concentrations of recombinant human BMP-7 for 72 hours before lysis and

SDS-PAGE



Preparation	
Shipping Temperature	ambient temperature
Formulation	10mM Acetic Acid
Reconstitution	Briefly centrifuge the vial before opening. It is recommended to reconstitute the protein in 10mM Acetic Acid containing 0.1% endotoxin-free recombinant human serum albumin (HSA).

Stability and Storage	Product Form	Temperature Conditions	Storage Time (From Date of Receipt)
	Lyophilized	-20°C to -80°C	Until Expiry Date
	Lyophilized	Room Temperature	2 weeks
	Reconstituted as per CofA	-20°C to -80°C	6 months
	Reconstituted as per CofA	4°C	1 week
Avoid repeated freeze-thaw cycles.			

www.ptglab.com

Document #: FR-QA118-101 Rev 0
Data Sheet Version #: 1

Proteintech Group, Inc.
5500 Pearl Street, Suite 400 Rosemont, IL 60612
t: 1-888-478-4522; f: 1-312-455-8408
Email: proteintech@ptglab.com