

IHC*easy* STAT2 Ready-To-Use IHC Kit

Catalog Number: **KHC1622**

General Information

Sample type:
FFPE tissue

Cited sample type:

Reactivity:
Human, Mouse, Rat

Cited Reactivity:

Assay type:
Immunohistochemistry

Primary antibody type:
Rabbit Polyclonal

Secondary antibody type:
Polymer-HRP-Goat anti-Rabbit

Kit Component

Component	Size	Concentration
Antigen Retrieval Buffer	100 mL	50×
Washing Buffer	100 mL ×2	20×
Blocking Buffer	5 mL	RTU
Primary Antibody	5 mL	RTU
Secondary Antibody	5 mL	RTU
Chromogen Component A	0.2 mL	RTU
Chromogen Component B	4 mL	RTU
Signal Enhancer	5 mL	RTU
Counter Staining Reagent	5 mL	RTU
Mounting Media	5 mL	RTU
Control Slide	1 slide (Optional)	FFPE
Datasheet	1 Copy	
Manual	1 Copy	

Storage Instructions

All the reagents are stored at 2-8°C. The kit is stable for 6 months from the date of receipt.


Background

STAT2, also named as p113, belongs to the transcription factor STAT family. It is a signal transducer and activator of transcription that mediates signaling by type I IFNs (IFN-alpha and IFN-beta). Following type I IFN binding to cell surface receptors, Jak kinases (TYK2 and JAK1) are activated, leading to tyrosine phosphorylation of STAT1 and STAT2. The phosphorylated STATs dimerize and associate with ISGF3G/IRF-9 to form a complex termed ISGF3 transcription factor, that enters the nucleus. ISGF3 binds to the IFN stimulated response element (ISRE) to activate the transcription of IFN stimulated genes, which drive the cell in an antiviral state. It also interacts with CRSP2, CRSP6, Simian virus 5 protein V, rabies virus phosphoprotein, IFNAR1 and IFNAR2. Its interaction with dengue virus NS5 inhibits the phosphorylation of STAT2, and, when all viral proteins are present (polyprotein), STAT2 is targeted for degradation.


Synonyms

ISGF 3, P113, STAT113, STAT2


Selected Validation Data




Immunohistochemical analysis of paraffin-embedded human oesophagus cancer tissue slide using KHC1622 (STAT2 IHC Kit).




Immunohistochemical analysis of paraffin-embedded human urothelial carcinoma tissue slide using KHC1622 (STAT2 IHC Kit).




Immunohistochemical analysis of paraffin-embedded mouse bladder tissue slide using KHC1622 (STAT2 IHC Kit).




Immunohistochemical analysis of paraffin-embedded mouse lung tissue slide using KHC1622 (STAT2 IHC Kit).



Immunohistochemical analysis of paraffin-embedded rat spleen tissue slide using KHC1622 (STAT2 IHC Kit).



Immunohistochemical analysis of paraffin-embedded rat bladder tissue slide using KHC1622 (STAT2 IHC Kit).



Immunohistochemical analysis of paraffin-embedded rat lung tissue slide using KHC1622 (STAT2 IHC Kit).