



# IHCeasy SP3 Ready-To-Use IHC Kit

Catalog Number: KHC1669

General Information

Sample type: FFPE tissue Cited sample type: Reactivity: Human, Mouse Cited Reactivity: Assay type: Immunohistochemistry Primary antibody type: Rabbit Polyclonal

Secondary antibody type: Polymer-HRP-Goat anti-Rabbit

#### Kit Component

Antigen Retrieval Buffer 100 mL	50×
Washing Buffer 100 mL × 2	20×
Blocking Buffer 5 mL	RTU
Primary Antibody 5 mL	RTU
Secondary Antibody 5 mL	RTU
Chromogen Component A 0.2 mL	RTU
Chromogen Component B 4 mL	RTU
Signal Enhancer 5 mL	RTU
Counter Staining Reagent 5 mL	RTU
Mounting Media 5 mL	RTU
Control Slide 1 slide (Optional)	FFPE
Datasheet 1 Copy	
Manual 1 Copy	

### **Storage Instructions**

All the reagents are stored at 2-8°C. The kit is stable for 6 months from the date of receipt.


#### Background


SP3, also named as transcription factor Sp3, is a 781 amino acid protein, which contains 3 C2H2-type zinc fingers and belongs to the Sp1 C2H2-type zinc-finger protein family. SP3 localizes to the nuclear periphery and in nuclear dots when sumoylated. Some localization in PML nuclear bodies. SP3 is acetylated by histone acetyltransferase p300, deacetylated by HDACs and sumoylated on all isoforms. Sumoylated on 2 sites in longer isoforms with Lys-551 being the major site. Sumoylation at this site promotes nuclear localization to the nuclear periphery, nuclear dots and PML nuclear bodies. Sumoylation on Lys-551 represses the transactivation activity, except for the largest isoform, L-Sp3, which has little effect on transactivation. Alternate sumoylation and acetylation at Lys-551 also control transcriptional activity.

## Synonyms

SP3, Sp3 transcription factor, SPR 2, Transcription factor Sp3

# Selected Validation Data





Immunohistochemical analysis of paraffinembedded human urothelial carcinoma tissue slide using KHC1669 (SP3 IHC Kit).

Immunohistochemical analysis of paraffinembedded mouse lung tissue slide using KHC1669 (SP3 IHC Kit).