

For Research Use Only

Human CD3/CD28 T cell Activation Beads Kit

Catalog Number: KMS310



Description

T lymphocytes are activated through MHC class II molecules on antigen-presenting cells (APC). After activation, T cells expand rapidly and secrete various cytokines. The Human CD3/CD28 T cell Activation Beads Kit is designed to activate and expand human T cells. This Kit contains Biotinylated anti-human CD3 and CD28 antibodies, as well as cell culture grade Streptavidin magnetic beads. After Streptavidin magnetic beads are loaded with biotinylated CD3/CD28, they mimic antigen presenting cells and can activate resting T lymphocytes from human PBMCs or purified T lymphocytes. After 2-3 days of activation, magnetic beads can easily be removed by magnet. Further T lymphocytes expansion will require human cytokines for *in vitro* culture.

Components

KMS310-10:
· MS310A-10: 100µL 0.1mg/mL functional grade Biotin-CD3 (clone: OKT3)
· MS310B-10: 100µL 0.2mg/mL functional grade Biotin-CD28 (clone: 28.2)
· MS310C-10: 500µL cell culture grade Streptavidin magnetic beads (2×10^8 beads/mL)

KMS310-100:
· MS310A-100: 1mL 0.1mg/mL functional grade Biotin-CD3 (clone: OKT3)
· MS310B-100: 1mL 0.2mg/mL functional grade Biotin-CD28 (clone: 28.2)
· MS310C-100: 5mL cell culture grade Streptavidin magnetic beads (2×10^8 beads/mL)

Package

10test/100test

Storage

2-8°C

Storage buffer

Biotin antibody buffer: PBS and 0.5% HSA, pH7.4
Streptavidin bead buffer: PBS and 0.5% HSA, 2mM EDTA, pH7.4

Reactivity

Human


Beads diameter

2.7 µm


Recommend usage

10µL Biotin-CD3, 10µL Biotin-CD28 and 50µL Streptavidin magnetic beads; mix well and add 30µL buffer for incubation. 100µL antibody-loaded magnetic beads contains 1×10^7 magnetic particles and is used to activate 1×10^7 purified human T lymphocytes or human PBMCs in 2mL cell culture media (1:1 bead to cell ratio).

Results



Representative example of human PBMCs activation: 2.5×10^6 human PBMCs in 0.5mL cell culture media are activated by Human CD3/CD28 T Cell Activation Beads Kit for 3 days at a 1:1 bead to cell ratio. Unstimulated cells (left panel) and stimulated cells (right panel) are stained with fluorescence flow antibodies APC-CD69 and FITC-... CD25. Total viable lymphocytes are gated. The activation effect of the Human CD3/CD28 T Cell Activation Beads Kit was tested on human PBMCs from three different donors.



Representative example of isolated T lymphocytes activation: 2.5×10^6 human CD3+ lymphocytes purified by Human CD3 T Cell Isolation Kit (KMS309) in 0.5mL cell culture media are activated by Human CD3/CD28 T Cell Activation Beads Kit for 3 days at a 1:1 bead to cell ratio. Unstimulated cells (left panel) and stimulated cells (right panel) are... stained with fluorescence flow antibodies APC-CD69, PE-CD25 and PB-CD4. Total viable lymphocytes are gated. The activation effect of the Human CD3/CD28 T Cell Activation Beads Kit was tested on human PBMCs from three different donors.