For Research Use Only

## NGAL Recombinant Matched Antibody Pair, PBS Only

Proteintech®
Antibodies | ELISA kits | Proteins
Www.ptgcn.com

Conjugate:

Full name:

lipocalin 2

Conjugate:

Full name:

lipocalin 2

Gene ID:

3934

Unconjugated

Gene ID:

3934

Unconjugated

Catalog Number: MP00770-1

Capture Antibody Information Catalog Number: 82841-9-PBS Host:

Rabbit human

Isotype: GenBank:
IgG NP\_005555.2

Purification Method: Protein A purfication

Detection Antibody Information

Catalog Number: 82841-2-PBS Host: Rabbit Isotype:

Purification Method: Protein A purfication

IgG

Tested Ap

Tested Applications: Sandwich ELISA \_

Clone ID:

Reactivity:

Clone ID:

Reactivity:

GenBank:

NP\_005555.2

human

218

2N10

19.5-1250 pg/mL (Sandwich ELISA)

Recommended Dilutions:

It is recommended that this reagent should be titrated in each testing system to obtain optimal results.

**Product Information** 

**Applications** 

 $MP00770\text{-}1\,targets\,NGAL\,in\,immuno assays\,as\,a\,matched\,antibody\,pair.\,Validated\,in\,Sandwich\,ELISA.$ 

Capture antibody: NGAL Recombinant antibody, PBS Only (Capture) 82841-9-PBS (2N10). 100  $\,\mu$  g. Concentration 1 mgl/ml.

Detection antibody: NGAL Recombinant antibody, PBS Only (Detector) 82841-2-PBS (2L8). 100  $\,\mu$  g. Concentration 1 mgl/ml.

Alternative NGAL matched antibody pairs: MP00302-1, MP00302-2, MP00302-3, MP00302-4

Unconjugated rabbit recombinant monoclonal antibody pair in PBS only storage buffer at a concentration of 1 mg/mL, ready for conjugation. Created using Proteintech's proprietary in-house recombinant technology. Recombinant production enables unrivalled batch-to-batch consistency, easy scale-up, and future security of supply.

Matched antibody pairs are designed for use in a variety of assays and platforms that require matched antibody pairs.

Antibody use should be optimized for each application and assay.

Storage

Storage:


Store at -80°C.


The product is shipped with ice packs. Upon receipt, store it immediately at -80°C


Storage buffer:

PBS only


## **Selected Validation Data**









Plasma of fifteen individual healthy human donors was measured. The NGAL concentration of detected samples was determined to be 38,260.8 pg/mL with a range of 17,778.7 - 90,593.8 pg/mL.



Human peripheral blood mononuclear cells (PBMC) were cultured unstimulated or stimulated with 10  $\,\mu$  g/ml PHA for 1 days. The mean NGAL concentration was determined to be 18,080.8 pg/mL in unstimulated PBMC supernatant, 32,268.8 pg/mL in PHA stimulated PBMC supernatant.



Saliva of eight individual healthy human donors was measured. The NGAL concentration of detected samples was determined to be 158,732.5 pg/mL with a range of 52,136.9 - 388,400.2 pg/mL



Urine of eight individual healthy human donors was measured. The NGAL concentration of detected samples was determined to be 11,654.6 pg/mL with a range of 1,964.2 - 21,150.4 pg/mL