For Research Use Only

Clusterin Recombinant Matched Antibody Pair, PBS Only



Catalog Number: MP00986-3

Capture Antibody Information

Detection Antibody

Catalog Number: 84067-1-PBS Host: Rabbit

Isotype:

Purification Method:

human

Protein A purification

Catalog Number: 84067-3-PBS

Host: Rabbit Isotype: IgG

Purification Method: Protein A purification

Applications

Information

Tested Applications:

Sandwich ELISA

Clone ID: 241277A4

Reactivity: GenBank: BC010514

Clone ID: 241277H2

Reactivity: human GenBank: BC010514

Recommended Dilutions:

Conjugate:

Full name:

clusterin

Gene ID:

Conjugate:

Full name:

clusterin

Gene ID:

1191

Unconjugated

1191

Unconjugated

It is recommended that this reagent should be titrated in each testing system to obtain optimal results.

Product Information

MP00986-3 targets Clusterin in immunoassays as a matched antibody pair. Validated in Sandwich ELISA.

97.7-6250 pg/mL (Sandwich ELISA)

Capture antibody: Clusterin Recombinant antibody, PBS Only (Capture) 84067-1-PBS (241277A4). 100 µg.

Detection antibody: Clusterin Recombinant antibody, PBS Only (Detector) 84067-3-PBS (241277H2). 100 $\,\mu$ g. Concentration 1 mgl/ml.

Unconjugated rabbit recombinant monoclonal antibody pair in PBS only storage buffer at a concentration of 1 mg/mL, ready for conjugation. Created using Proteintech's proprietary in-house recombinant technology. Recombinant production enables unrivalled batch-to-batch consistency, easy scale-up, and future security of supply.

Matched antibody pairs are designed for use in a variety of assays and platforms that require matched antibody pairs.

Antibody use should be optimized for each application and assay.

Storage


Storage:

Store at -80°C.


The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

PBS only


Selected Validation Data




Sandwich ELISA standard curve of MP00986-3, Human Clusterin Recombinant Matched Antibody Pair - PBS only. 84067-1-PBS was coated to a plate as the capture antibody and incubated with serial dilutions of standard Eg0225. 84067-3-PBS was HRP conjugated as the detection antibody. Range: 97.7-6250 pg/mL




Serum of six individual healthy human donors was measured. The human Clusterin concentration of detected samples was determined to be 31,152.14 ng/mL with a range of 15,052.59 - 51,157.85 ng/mL



Saliva of four individual healthy human donors was measured. The human Clusterin concentration of detected samples was determined to be 152.81 ng/mL with a range of 95.56 - 231.47 ng/mL



Urine of four individual healthy human donors was measured. The human Clusterin concentration of detected samples was determined to be 40.81 ng/mL with a range of 16.72 - 69.41 ng/mL



HepG2 were cultured in DMEM supplemented with 10% fetal bonive serum, 2.5 mM L-glutamine, 100 U/mL penicillin, and 100 μ g/mL streptomycin sulfate. An aliquot of the cell culture supernate was removed, assayed for human Clusterin, and measured 140.90 ng/mL