

Human CD19 Magnetic Beads

Catalog Number: MS005

Description

CD19 is a 95 kDa, type I transmembrane glycoprotein that belongs to the immunoglobulin superfamily. It is expressed by B cells and follicular dendritic cells and regulates B cell lineage commitment during hematopoietic stem cell differentiation. It is observed that 6%-23% human PBMC are CD19 positive. Human CD19 Magnetic Beads can be used for isolation or depletion of human CD19 B lymphocytes from PBMC, whole blood, or other sample types. Following incubation with human CD19 antibody conjugated magnetic beads, the cell sample is placed on a magnet. CD19+ cells remain attached to magnetic beads after separation and can be used for downstream applications, such as in expansion of cells, but are not suitable for flow cytometry analysis. CD19- cells remain in supernatant and could be used for further applications.

Components

MS005-10: 100 µL 10 mg/mL Human CD19 Magnetic Beads
 MS005-100: 1 mL 10 mg/mL Human CD19 Magnetic Beads

Package

10 test/100 test

Storage

2-8°C

Storage buffer

PBS, pH7.4, 0.2% BSA and 0.05% Sodium Azide

Reactivity

Human


Recommend usage

10 µL Human CD19 Magnetic Beads for 1×10^7 cells

Beads Diameter 2.7 µm

Results

Representative example of enrichment and depletion



Following depletion of CD19+ cells, supernatant cell suspension was stained with PB450-CD3(clone: HIT3a) and CL647-CD19(clone: SJ25C1) antibodies. CD45 positive cells are gated in the analysis. Left panel: CD3-CD19+ cells before selection. Right panel: CD3-CD19+ cells after depletion. Human CD19 Magnetic Beads are tested using PBMC from three different donors.

For technical support and original validation data for this product please contact:

T: 027-87531629 E: Proteintech-CN@ptglab.com W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.