For Research Use Only

PE Anti-Mouse TCR Beta (H57-597)

Catalog Number:PE-65106




Catalog Number: PE-65106 Size: 100ug, 0.2 mg/ml	GenBank Accession Number: GeneID (NCBI): 21577 Full Name: T-cell receptor beta chain	Purification Method: Affinity purification CloneNo.: H57-597 Excitation/Emission maxima wavelengths: 496 nm, 565 nm / 578 nm
Isotype: IgG		
Tested Applications: FC		
Species Specificity: mouse		
homology with immunoglobulins (is expressed on alpha/beta TCR-be associated with CD3. The TCR/CD3 gamma/delta heterodimer coexpre epsilon, zeta, and eta (PMID: 18262 (MHC) molecules. TCR-mediated po	PMID: 2144901). TCR beta is a memb aring T cells and thymocytes. The TC complex of T-lymphocytes consists essed at the cell surface with the inva 55). The TCR recognizes antigens bou eptide-MHC recognition is transmitte	er of the immunoglobulin superfamily and R heterodimers are noncovalently of either a TCR alpha/beta or TCR ariant subunits of CD3 labeled gamma, delta, und to major histocompatibility complex
Storage Buffer:		
	PE-65106 Size: 100ug, 0.2 mg/ml Source: Armenian Hamster Isotype: IgG Tested Applications: FC Species Specificity: mouse T-cell receptor (TCR) consists of her homology with immunoglobulins (is expressed on alpha/beta TCR-be associated with CD3. The TCR/CD3 gamma/delta heterodimer coexpre epsilon, zeta, and eta (PMID: 18262 (MHC) molecules. TCR-mediated pr intracellular signal transduction (P Storage: Stora at 2-8°C. Avoid exposure to li Storage Buffer:	PE-65106 GeneID (NCBI): Size: 21577 100ug, 0.2 mg/ml Full Name: Source: T-cell receptor beta chain Armenian Hamster Isotype: IgG IgG Tested Applications: FC Species Specificity: mouse T-cell receptor (TCR) consists of heterodimeric glycoproteins (TCR alpha homology with immunoglobulins (PMID: 2144901). TCR beta is a memb is expressed on alpha/beta TCR-bearing T cells and thymocytes. The TC associated with CD3. The TCR/CD3 complex of T-lymphocytes consists.gamma/delta heterodimer coexpressed at the cell surface with the inva epsilon, zeta, and eta (PMID: 1826255). The TCR recognizes antigens bot (MHC) molecules. TCR-mediated peptide-MHC recognition is transmitte intracellular signal transduction (PMID: 11985657). Storage: Storage: Storage: Storage:

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1x10^{^6} mouse splenocytes were surface stained with 0.25 ug PE Anti-Mouse TCR Beta (H57-597) (PE-65106, Clone: H57-597) or 0.25 ug PE Armenian Hamster IgG Isotype Control (PIP) (PE-65210, Clone: PIP). Cells were not fixed.