For Research Use Only

PE Anti-Mouse CD8a (5H10-1)

Catalog Number: PE-65205




Basic Information	Catalog Number: PE-65205	GenBank Accession Number: BC030679	Purification Method: Affinity purification
	Size: 100ug, 0.2 mg/ml	GenelD (NCBI): 12525	CloneNo.: 5H10-1
	Source: Rat Isotype:	UNIPROT ID: P01731 Full Name: CD8 antigen, alpha chain	Excitation/Emission maxima wavelengths: 496 nm, 565 nm / 578 nm
	IgG2b, lambda		
Applications	Tested Applications: FC		
	Species Specificity: Mouse		
Background Information	CD8 is a transmembrane glycoprotein composed of two disulfide-linked chains. It can be present as a homodimer of CD8 α or as a heterodimer of CD8 α and CD8 β (PMID: 3264320; 8253791). CD8 is found on most thymocytes. The majority of class I-restricted T cells express mostly the CD8 α β heterodimer while CD8 α α homodimers alone have been found on some gut intraepithelial T cells, on some T cell receptor (TCR) γ δ T cells and on NK cells (PMID: 2111591; 1831127; 8420975). CD8 acts as a co-receptor that binds to MHC class-I and participates in cytotoxic T cell activation (PMID: 8499079). During T cell development, CD8 is required for positive selection of CD4-/CD8+T cells (PMID: 1968084).		
Storage	Storage: Store at 2-8°C. Avoid exposure to li Storage Buffer: PBS with 0.1% sodium azide and 0	ight. Stable for one year after shipmer .5% BSA.	nt.

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1X10^{^6} C 57BL/6 mouse splenocytes were surface co-stained with CoraLite® Plus 647 Anti-Mouse CD3 (17A2) and 0.1 ug PE Anti-Mouse CD8a (PE-65205, Clone: 5H10-1) or 0.1 ug PE Rat IgG2b Isotype Control (LTF-2). Cells were not fixed.