## For Research Use Only

## PE Anti-Human CD107a / LAMP1 (H4A3) Mouse IgG2a Recombinant Antibody



Catalog Number: PE-65596

**Basic Information** 

Catalog Number: PE-65596

Concentration: 100tests, 5 ul/test

Source: Mouse Isotype:

IgG2a

GenBank Accession Number:

BC006345 GenelD (NCBI): 3916

ENSEMBL Gene ID: ENSG00000185896 UNIPROT ID:

P11279 Full Name:

lysosomal-associated membrane

protein 1 Calculated MW: 45 kDa Purification Method:

Protein A purification

CloneNo.: H4A3

**Recommended Dilutions:** 

FC (Intra): 5 ul per 10^6 cells in a 100  $\,$ 

μl suspension

Excitation/Emission maxima

wavelengths:

496 nm, 565 nm / 578 nm

**Applications** 

Tested Applications: FC (Intra)

Species Specificity:

human

Positive Controls:

FC (Intra): Jurkat cells,

## **Background Information**

LAMP1 (CD107a) is a heavily glycosylated membrane protein enriched in the lysosomal membrane. LAMP1 is extensively glycosylated with asparagine-linked oligosaccharides which protect it from intracellular proteolysis (PMID: 10521503). Although LAMP1 is expressed largely in the endosome-lysosomal membrane of cells, it is also found on the plasma membrane (PMID: 16168398). Elevated LAMP1 expression at the cell surface has also been detected during platelet and granulocytic cell activation, as well as in some tumor cells (PMID: 29085473). LAMP1 functions to provide selectins with carbohydrate ligands. This protein has also been shown to be a marker of degranulation on lymphocytes such as CD8+ and NK cells and may also play a role in tumor cell differentiation and metastasis (PMID: 18835598; 29085473; 9426697).


Storage

Storage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 0.09% sodium azide and 0.5% BSA, pH7.3

## Selected Validation Data



1x10^6 Jurkat cells were intracellularly stained with 5 ul PE Anti-Human CD107a / LAMP1 (H4A3) Mouse IgG2a RecAb (PE-65596, Clone: H4A3) (red) or PE Mouse IgG2a Isotype Control (C1.18.4) (PE-65208, Clone: C1.18.4) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer.