For Research Use Only

## PE Anti-Mouse CD206 Rabbit Recombinant Antibody

Catalog Number: PE-98031



**Purification Method:** 

Protein A purification

**Basic Information** 

Catalog Number: GenBank Accession Number: PE-98031 NM\_008625.2

Concentration: Genel D (NCBI): CloneNo.: 100ug, 200  $\mu$  g/ml 17533 240344D12 Source: UNIPROT ID: Recommended Dilutions:

Rabbit Q2HZ94 FC: 0.1 ug per 10^6 cells in 100  $\,\mu$  l sotype: Full Name: suspension

G mannose receptor, C type 1 Excitation/Emission maxima

 Calculated MW:
 wavelengths:

 165 kDa
 496 nm, 565 nm / 578 nm

**Applications** 

Tested Applications:

Species Specificity:

mouse

Positive Controls:

FC: mouse peritoneal macrophages,

## **Background Information**

CD206, also named as MMR, CLEC13D and MRC1, is a type I membrane receptor that mediates the endocytosis of glycoproteins by macrophages. CD206 has been shown to bind high-mannose structures on the surface of potentially pathogenic viruses, bacteria, and fungi so that they can be neutralized by phagocytic engulfment. CD206 is a 170 kDa transmembrane protein which contains 5 domains: an amino-terminal cysteine-rich region, a fibronectin type II repeat, a series of eight tandem lectin-like carbohydrate recognition domains (responsible for the recognition of mannose and fucose), a transmembrane domain, and an intracellular carboxy-terminal tail. It is expressed on most tissue macrophages, in vitro derived dendritic cells, lymphatic and sinusoidal endothelia.


Storage

Storage:


Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer:

PBS with 0.09% sodium azide and 0.5% BSA, pH7.3

## **Selected Validation Data**



1x10^6 mouse peritoneal macrophages were surface stained with APC Anti-Mouse CD107b / LAMP2, and 0.1 ug PE Anti-Mouse CD206 Rabbit RecAb (PE-98031, Clone:240344D12) or 0.1 ug PE Rabbit IgG Isotype Control RecAb (PE-98136, Clone: 240953C9). Cells were incubated with FC Receptor Block prior to staining and fixing. Cells were not fixed.



1x10^6 mouse peritoneal macrophages were surface stained with 0.1 ug PE Anti-Mouse CD206 Rabbit RecAb (PE-98031, Clone: 240344D12) (red) or 0.1 ug PE Rabbit 1gG 1sotype Control RecAb (PE-98136, Clone: 240953C9) (blue). Cells were incubated with FC Receptor Block prior to staining and fixing. Cells were not fixed.