## For Research Use Only

## FcZero-rAb™ PE Anti-Human CD8a (OKT8) Rabbit IgG Recombinant Antibody



Catalog Number: PE-FcA65506

**Basic Information** 

Catalog Number:

PE-FcA65506

Size:

100tests, 5 ul/test

Source: Rabbit

Isotype:

GenBank Accession Number:

BC025715 GeneID (NCBI):

ENSEMBL Gene ID:

ENSG00000153563

Full Name: CD8a molecule

Calculated MW: 235 aa, 26 kDa

**Purification Method:** 

Protein A purification

CloneNo.: ОКТ8

Excitation/Emission maxima wavelengths:

496 nm, 565 nm / 578 nm

**Applications** 

**Tested Applications:** 

Species Specificity:

human

## **Background Information**

CD8 is a transmembrane glycoprotein composed of two disulfide-linked chains. It can be present as a homodimer of CD8  $\alpha$  or as a heterodimer of CD8  $\alpha$  and CD8  $\beta$  (PMID: 3264320; 8253791). CD8 is found on most thymocytes. The majority of class I-restricted T cells express mostly the CD8  $\alpha$   $\beta$  heterodimer while CD8  $\alpha$   $\alpha$  homodimers alone have been found on some gut intraepithelial T cells, on some T cell receptor (TCR)  $\gamma$   $\delta$  T cells and on NK cells (PMID: 2111591; 1831127; 8420975). CD8 acts as a co-receptor that binds to MHC class-I and participates in cytotoxic T-cell activation (PMID: 8499079). During T cell development, CD8 is required for positive selection of CD4-/CD8+T cells (PMID: 1968084).


Storage

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:

PBS with 0.09% sodium azide and 0.5% BSA.

## Selected Validation Data



1x10^6 human PBMCs were surface stained with APC Anti-Human CD4 and 5 ul PE Anti-Human CD8a (OKT8) Rabbit IgG Recombinant Antibody (PE-FcA65506, Clone: OKT8) or PE Rabbit IgG Isotype Control Recombinant Antibody (PE-FcA98136, Clone: 240953C9). Cells were not fixed.