

Catalog Number: atd

1 Publications

Basic Information

Catalog Number:

atd

Applications:

IP, Co-IP

Conjugate:

Magnetic Particles M-270, size: 2.8 µm

Host:

Alpaca

Type:

Nanobody

Class:

Recombinant

Description

The ChromoTek HA-Trap Magnetic Particles M-270 consists of an anti-HA-tag Nanobody/VHH, which is coupled to magnetic particles. It can be used for the immunoprecipitation of HA-tagged proteins from cell extracts of various organisms such as humans, mice, dogs, plants, and yeast. It is highly recommended when very large proteins/complexes are being investigated.

Specificity/Target

Binds specifically to the HA-tag (sequence YPYDVPDYA) fused to a protein of interest at N-, C- or internal position. Please note that the affinity is highest for a C-terminal fusion. There is no cross-reactivity to other common peptide tags such as the His6-tag, FLAG-tag, Spot-Tag, V5-tag, Strep-tag, or C-tag (other tags not tested). Background binding to host cell proteins from a range of organisms such as human, mouse and dog cell lines or yeast and plants is low.

Elution buffer

2x SDS-sample buffer (Lamli)

Affinity (K_D)

6 nM for C-terminal HA-tags and ca. 180 nM for N-terminal fusions.

Storage


Storage:

Shipped at ambient temperature. Upon receipt store at +4°C. Stable for one year. DO not freeze!


Storage Buffer:

PBS with 0.09% sodium azide


Selected Validation Data




The HA-Trap Magnetic Particles M-270 (left) and a competitor resin (right) were used to immunoprecipitate TOM70-HA fusion protein from either untransfected (mock) HEK293T cells or HEK293T cell transfected with full-length TOM70-HA construct. Immunoprecipitation with HA-Trap Magnetic Particles M-270 results in cleaner, single-band pull-downs without any heavy and light chain contamination. SDS-PAGE analysis was done on sample: (B) fract



The HA-Trap Magnetic Particles M-270 was used to immunoprecipitate HA-PCNA fusion protein from HEK293T cells. HA-PCNA protein was released from the trap through a two-step competitive elution utilizing HA-peptide (ap). Samples from the Input (I), Flow-Through (F), 1st elution (E1), 2nd elution (E2), and residual (R) fractions were analyzed through WB. PCNA Monoclonal Antibody (60097-1-Ig) and Multi-rAb HRP-Goat Anti-Mouse Recombinant Secondary Antibody (RGAM001) were used in the WB analysis. Note: PCNA forms trimers resulting in co-elution of endogenous PCNA proteins with HA-tagged PCNA.



WB detection of TOM70-HA fusion protein following immunoprecipitation with HA-Trap Magnetic Particles M-270 from either untransfected (mock) HEK293T cells or HEK293T cells transfected with full-length TOM70-HA construct. Samples from the Input (I), Flow-Through (F), and Bound (B) fractions were used in the WB analysis. Detection was completed using TOM70 Monoclonal Antibody (66593-1-Ig) and Multi-rAb HRP-Goat Anti-Mouse Recombinant Secondary Antibody (RGAM001).



The HA-Trap Magnetic Particles M-270 was used to immunoprecipitate HA-PCNA and PCNA-HA proteins from transfected HEK293T cells. WB analysis was done on samples from the Input (I), Flow-Through (F), and Bound (B) fractions of the IP using PCNA Monoclonal Antibody (60097-1-Ig) and Multi-rAb HRP-Goat Anti-Mouse Recombinant Secondary Antibody (RGAM001). The HA-Trap is successful in pulling down HA-tagged PCNA regardless of whether the tag is fused to the N- or C-terminal. Note: PCNA forms trimers, resulting in co-elution of endogenous PCNA with HA-tagged PCNA.