

For Research Use Only

His Fab-Trap® Agarose



www.ptgcn.com

Catalog Number: hfa

Basic Information

Catalog Number:

hfa

Applications:

IP, Co-IP

Conjugate:

Agarose beads; ~90 µm (cross-linked 4% agarose beads)

Host:

Mouse

Type:

Fab-fragment

Class:

Recombinant

Description

The ChromoTek His Fab-Trap® Agarose consists of an anti-His-tag Fab fragment, which is covalently bound to agarose beads. It can be used for the immunoprecipitation of proteins fused to the His-tag from cell extracts of various organisms.

Specificity/Target

Binds specifically to the His-tag (also known as hexahistidine or His6-tag) fused to a protein of interest at N-, C- or internal position. Please note that the affinity is highest for a C-terminal fusion (dissociation constant K_D of 10 nM for a C-terminal His-tag and 220 nM for an N-terminal His-tag).

Elution buffer

2x SDS-sample buffer (Lämmli), 200 mM glycine pH 2.5 or 100 µM His-Peptide

Affinity (K_D)

10 nM for a C-terminal His-tag and 220 nM for a N-terminal His-tag

Storage

Storage:

Shipped at ambient temperature. Upon receipt store at +4°C. Stable for one year. Do not freeze!

Storage Buffer:

PBS pH 7.4. Preservative: 0.09% sodium azide

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: www.ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




IP using His Fab-Trap® Agarose (hfa) from untransfected HEK293T (mock) and HEK293T cells transfected with either His-TOM70 or TOM70-His fusion proteins. WB detection of TOM70 protein was also done on samples from the Input (I), Flow-Through (F), and Bound (B) fractions using Anti-TOM70 Polyclonal Antibody (14528-1-AP) and ChromoTek Nano-Secondary alpaca anti-rabbit IgG, CoraLite Plus 488 (srb2GCL488-1).



IP of His-TOM70 and TOM70-His proteins using either His Fab-Trap Agarose (left) or a competitor resin product (right) from transfected HEK293T cells. The His Fab-Trap pulls down the TOM70 proteins more efficiently and with less background when compared to the competitor.



His Fab-Trap® Agarose (hfa) was used for the Co-IP of AurKinB-His fusion protein and interacting partners from untransfected (mock) and transfected HEK293T cells. Multiplexed WB was done on Input (I), Flow-Through (F), and Bound (B) fractions of the IP. Detection was completed using His-tag Monoclonal Antibody (66005-1-Ig) labeled with FlexAble CoraLite Plus 647 Kit for Mouse IgG (KFA023), CDC37 Monoclonal Antibody (66420-1-Ig) labeled with FlexAble CoraLite Plus 750 Kit for Mouse IgG2a (KFA044), and HSP90AB1 Monoclonal Antibody (67450-1-Ig) labeled with FlexAble CoraLite Plus 488 Kit for Mouse IgG2b (KFA061).



IP of GFP-His fusion protein from transfected HEK293T cells using His Fab-Trap Agarose (hfa) followed by a two-step elution using His-Peptide (hp). I: Input, F: Flow-through, E1: 1st elution, E2: 2nd elution, R: Residual.