

Catalog Number: yt2a

## Basic Information

**Catalog Number:**  
yt2a

**Applications:**  
IP, Co-IP

**Conjugate:**  
Agarose beads; ~90 um (cross-linked 4% agarose beads)

**Host:**  
Alpaca

**Type:**  
Nanobody

**Class:**  
Recombinant

## Description

The ChromoTek Myc-Trap® 2.0 Agarose consists of an anti-Myc-tag VHH, which is coupled to agarose beads. It can be used for the immunoprecipitation of Myc-tag-fusion proteins from cell extracts of various organisms.

## Specificity/Target

Binds specifically to the Myc-tag (sequence EQKLISEEDL) at the N-terminus, C-terminus, or internal site of the fusion protein. Endogenous c-myc is NOT bound.

## Elution buffer

2x SDS-sample buffer (Lämmli), 200 mM glycine pH 2.5, 0.1 mg/ml ChromoTek 2x Myc-peptide (2yp) in PBS pH 7.4

## Affinity ( $K_D$ )


770 nM

## Storage


**Storage:**  
Shipped at ambient temperature. Upon receipt store at +4°C. Stable for one year. Do not freeze!

**Storage Buffer:**  
20% ethanol

## Selected Validation Data



Immunoprecipitation of GFP-Myc fusion protein from HEK293T cells using Myc-Trap® 2.0 Agarose (yt2a). IP was done using both un-transfected (mock) and transfected (GFP-myc) cells. I: Input, F: Flow-through, B: Bound.



Comparison of pull-down efficacy between the Myc-Trap® 2.0 Agarose (yt2a) (left) and the original Myc-Trap Agarose (yta) (right). Both products were used to immunoprecipitate GFP-myc fusion proteins from untransfected (mock) and transfected (GFP-myc) HEK293T cells. The Myc-Trap 2.0 has a higher binding capacity and is able to pull down more GFP-Myc protein than the Myc-Trap.